


Treatment-Naive Genotype 1a Without Cirrhosis

Recommended regimens listed by evidence level and alphabetically for:

Treatment-Naive Genotype 1a Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg) for patients without baseline NS5A RASs for elbasvir ^a	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^b	8 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
<hr/>		
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks	I, B
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
<p>^a Includes genotype 1a resistance-associated substitutions (RASs) at amino acid positions 28, 30, 31, or 93 known to confer antiviral resistance. If 1 or more RASs are present, another recommended regimen should be used.</p> <p>^b Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.</p>		

Last update: November 6, 2019