


Treatment-Naive Genotype 1a With Compensated Cirrhosis

Recommended regimens listed by evidence level and alphabetically for:

Treatment-Naive Genotype 1a Patients With Compensated Cirrhosis^a

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg) for patients without baseline NS5A RASs ^b for elbasvir	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) ^c	8 weeks	I, B

^a For [decompensated cirrhosis](#), please refer to the appropriate section.

^b Includes genotype 1a RASs at amino acid position 28, 30, 31, or 93 known to confer antiviral resistance. If 1 or more RASs are present, another recommended regimen should be used.

^c Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information. For patients with HIV/HCV coinfection, a treatment duration of 12 weeks is recommended.

Last update: August 27, 2020