


## Treatment-Naive Genotype 1b With Compensated Cirrhosis

Recommended regimens listed by evidence level and alphabetically for:

### Treatment-Naive Genotype 1b Patients With Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks <sup>c</sup>	I, B

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.

<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

<sup>c</sup> For HIV/HCV-coinfected patients, a treatment duration of 12 weeks is recommended.

For genotype 1b-infected, treatment-naive patients with compensated cirrhosis, there are 4 recommended regimens with comparable efficacy. The alternative regimen is classified as such because, compared to the recommended regimens, it requires a longer duration of treatment, involves greater prescribing complexity, is potentially less efficacious, and/or there are limited supporting data.

## Recommended Regimens

### Elbasvir/Grazoprevir

The recommendation for use of daily fixed-dose elbasvir (50 mg)/grazoprevir (100 mg) in cirrhotic patients with genotype 1 infection is based on 92 patients (22% of the study cohort) in the phase 3 C-EDGE trial who had Metavir F4 disease ([Zeuzem, 2015f](#)). SVR12 was 97% in the subgroup of cirrhotic patients. A similar 97% (28/29) SVR12 rate had previously been demonstrated in genotype 1 cirrhotic treatment-naive patients treated with 12 weeks of elbasvir/grazoprevir without ribavirin in the open-label phase 2 C-WORTHY trial, which enrolled both HCV-monoinfected and HIV/HCV-coinfected patients ([Lawitz, 2015c](#)). Presence or absence of cirrhosis does not appear to alter the efficacy of the elbasvir/grazoprevir regimen ([Lawitz, 2015c](#)); ([Zeuzem, 2017](#)).

### Glecaprevir/Pibrentasvir

EXPEDITION-1 investigated use of the daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) administered as three 100 mg/40 mg fixed-dose combination pills in DAA-naive (75%) or -experienced (interferon or peginterferon ± ribavirin, or sofosbuvir plus ribavirin ± peginterferon) patients with compensated cirrhosis. Of 146 patients with genotype 1, 2, 4, 5, or 6 given 12 weeks of glecaprevir/pibrentasvir, 145 (99%) achieved SVR12; all genotype 1b patients achieved SVR ([Forns, 2017](#)).

EXPEDITION-2, a study of glecaprevir/pibrentasvir in 153 HIV/HCV-coinfected adults with genotype 1, 2, 3, 4, 5, or 6, utilized 8 weeks of treatment for noncirrhotic patients and 12 weeks for cirrhotic patients (the recommended durations

approved by the FDA). The overall SVR12 rate was 98% and there were no observed virologic failures among the 94 patients with genotype 1 infection ([Rockstroh, 2017](#)). In EXPEDITION-1 and EXPEDITION-2, neither subtype (1a vs 1b) nor the presence of baseline RASs impacted SVR12 results in DAA-naive genotype 1 patients.

EXPEDITION-8 evaluated glecaprevir/pibrentasvir for a reduced duration of 8 weeks in 280 treatment-naive patients with compensated cirrhosis and genotype 1 (n=136, genotype 1b), 2, 4, 5 or 6 infection. Patients with a prior history of decompensation, hepatocellular carcinoma, and HIV or HBV coinfection were excluded from this study. SVR12 was 99% with no virologic failures ([Brown, 2018](#)).

## Ledipasvir/Sofosbuvir

The daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) was approved by the FDA for the treatment of genotype 1 infection in treatment-naive patients based on 2 registration trials: ION-1 (865 treatment-naive patients; those with cirrhosis were included) and ION-3 (647 treatment-naive patients; those with cirrhosis were excluded). ION-1 investigated length of treatment (12 weeks vs 24 weeks) and the need for ribavirin ([Afdhal, 2014a](#)). SVR12 rates were 97% to 99% across all study arms with no difference in SVR based on length of treatment, use of ribavirin, or genotype 1 subtype. Sixteen percent of participants enrolled were classified as cirrhotic. There was no difference in SVR12 rate in cirrhotic (97%) versus noncirrhotic patients (98%).

## Sofosbuvir/Velpatasvir

The daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) for 12 weeks was approved by the FDA for the treatment of genotype 1 infection in treatment-naive patients based on ASTRAL-1. This placebo-controlled trial involved a 12-week course of sofosbuvir/velpatasvir administered to 624 participants with genotype 1, 2, 4, 5, or 6 who were treatment-naive (n=423) or previously treated with interferon-based therapy, with or without ribavirin or a protease inhibitor (n=201); ([Feld, 2015](#)). Of the 328 genotype 1 patients included, 323 achieved SVR12 with no difference in SVR12 observed by subtype (98% 1a, 99% 1b). Of 121 participants (all genotypes) classified as having cirrhosis, 120 achieved SVR12 (99%). Baseline NS5A RASs (at 15% cutoff)—reported in 11% of genotype 1a and 18% of genotype 1b participant samples tested—did not influence SVR12 rate for genotype 1 ([Hézode, 2018](#)). Of the 2 virologic failures in ASTRAL-1 (<1% of treated participants), both were genotype 1 and had baseline RASs. There was no significant difference in the rates of adverse events in the sofosbuvir/velpatasvir vs placebo groups.

The phase 3 POLARIS-2 study randomized 941 DAA-naive patients with genotype 1, 2, 3, 4, 5, or 6 infection—19% of whom had compensated cirrhosis—to receive either 8 weeks of sofosbuvir (400 mg)/velpatasvir (100 mg)/voxilaprevir (100mg) or 12 weeks of sofosbuvir/velpatasvir ([Jacobson, 2017](#)). Of participants treated with sofosbuvir/velpatasvir, 170/172 (99%) with genotype 1a and 57/59 (97%) with genotype 1b achieved SVR with a single relapse observed with each subtype.

**Last update:** November 6, 2019

## Related References

Afdhal NH, Zeuzem S, Kwo PY, et al. [Ledipasvir and sofosbuvir for untreated HCV genotype 1 infection](#). *N Engl J Med*. 2014;370(20):1889-1898.

Brown RS, Hezode C, Wang S, et al. [Preliminary efficacy and safety of 8-week glecaprevir/pibrentasvir in patients with HCV genotype 1-6 infection and compensated cirrhosis: the EXPEDITION-8 study \[Abstract LB-7\]](#). *The Liver Meeting*. 2018.

Feld JJ, Jacobson IM, Hézode C, et al. [Sofosbuvir and velpatasvir for HCV genotype 1, 2, 4, 5, and 6 infection](#). *N Engl J Med*. 2015;373(27):2599-2607.

Forns X, Lee SS, Valdes J, et al. [Glecaprevir plus pibrentasvir for chronic hepatitis C virus genotype 1, 2, 4, 5, or 6 infection in adults with compensated cirrhosis \(EXPEDITION-1\): a single-arm, open-label, multicentre phase 3 trial.](#) *Lancet Infect Dis.* 2017;17(10):1062-1068.

Hezode C, Reau N, Svarovskaia ES, et al. [Resistance analysis in patients with genotype 1-6 HCV infection treated with sofosbuvir/velpatasvir in the phase III studies.](#) *J Hepatol.* 2018;68(5):895-903.

Jacobson IM, Lawitz E, Gane EJ, et al. [Efficacy of 8 weeks of sofosbuvir, velpatasvir, and voxilaprevir in patients with chronic HCV infection: 2 phase 3 randomized trials.](#) *Gastroenterology.* 2017;153(1):113-122.

Lawitz EJ, Gane EJ, Pearlman B, et al. [Efficacy and safety of 12 wks vs 18 wks of treatment w/ GRZ and ELB w/ or without RBV for HCV GT1 infection in previously untreated pts w/ cirrhosis and pts w/ previous null response w/ or without cirrhosis \(C-WORTHY\), randomised, open-label phase 2 trial.](#) *Lancet.* 2015;385(9973):1075-1086.

Rockstroh J, Lacombe K, Viani RM, et al. [Efficacy and safety of Glecaprevir/Pibrentasvir in patients co-infected with hepatitis C virus and human immunodeficiency virus-1: the EXPEDITION-2 Study \[Abstract LBP-522\].](#) In: *The International Liver Congress. EASL.* The International Liver Congress. EASL.; 2017. Available at: [http://dx.doi.org/10.1016/S0168-8278\(17\)30467-1](http://dx.doi.org/10.1016/S0168-8278(17)30467-1).

Zeuzem S, Ghalib R, Reddy KR, et al. [Grazoprevir-Elbasvir Combination Therapy for Treatment-Naive Cirrhotic and Noncirrhotic Patients With Chronic Hepatitis C Virus Genotype 1, 4, or 6 Infection: A Randomized Trial.](#) *Ann Intern Med.* 2015;163(1):1-13.

Zeuzem S, Mizokami M, Pianko S, et al. [NS5A resistance-associated substitutions in patients with genotype 1 hepatitis C virus: prevalence and effect on treatment outcome.](#) *J Hepatol.* 2017;66(5):910-918.