



Table 3. Commonly Used Abbreviations

Abbreviation	Definition and Notes		
ACA	Patient Protection and Affordable Care Act		
AFP	alpha-fetoprotein		
ALT	alanine aminotransferase		
AMP	average manufacturer price		
Anti-HCV	HCV antibody		
APRI	AST-to-platelet ratio index		
AST	aspartate aminotransferase		
AUC	area under the curve		
AWP	average wholesale price ^a		
BOC	boceprevir		
CBC	complete blood count		
CDC	Centers for Disease Control and Prevention		
CEA	cost-effectiveness analysis		
СТР	Child-Turcotte-Pugh (see below)		
СҮР	cytochrome P450		
DAA	direct-acting antiviral		
eGFR	estimated glomerular filtration rate		
ESRD	end-stage renal disease		
FDA	US Food and Drug Administration		
GFR	glomerular filtration rate		
HBsAg	hepatitis B virus surface antigen		
HBV	hepatitis B virus		





Table 3. Commonly Used Abbreviations

Published on HCV Guidance (https://www.hcvguidelines.org)

Abbreviation	Definition and Notes		
HCC	hepatocellular carcinoma		
HCV	hepatitis C virus Hepatitis C virus and HCV refer to the virus. Hepatitis C and HCV infection or HCV disease refer to the disease entity.		
ICER	incremental cost-effectiveness ratio		
IDU	injection drug use or user		
INR	international normalized ratio		
MELD	model for end-stage liver disease		
MSM	men who have sex with men		
NASH	nonalcoholic steatohepatitis		
NAT	nucleic acid testing		
NIH	National Institutes of Health		
NS3	HCV nonstructural protein 3		
NS5A	HCV nonstructural protein 5A		
OATP	organic anion-transporting polypeptide		
PBM	pharmacy benefit manager		
PCR	polymerase chain reaction		
P-gp	P-glycoprotein		
PreP	preexposure prophylaxis		
PWID	people who inject drugs		
QALY	quality-adjusted life-year		
RAS	resistance-associated substitution		
RBC	red blood cell(s)		
RBV	ribavirin		
RGT	response-guided therapy		
sAg	surface antigen		
SMV	simeprevir		
SOF	sofosbuvir		
SOF	SofoSbuvir		





Table 3. Commonly Used Abbreviations

Published on HCV Guidance (https://www.hcvguidelines.org)

Abbrevietien	Definition and Notes			
Abbreviation	Definition and Notes			
SVR12 (or 24 or 48, etc)	sustained virologic response at 12 weeks (or at 24 weeks, or at 48 weeks, etc)			
TSH	thyroid-stimulating hormone			
TVR	telaprevir			
ULN	upper limit of normal			
USPSTF	US Preventive Services Task Force			
WAC	wholesale acquisition cost ^b			
^a "List price" for wholesale pharmacies to purchase drugs				

^b Typically, approximately 17% off of AWP

Child-Turcotte-Pugh (CTP) Classification of the Severity of Cirrhosis					
	CLASS A	CLASS B	CLASS C		
Total Points	5-6	7-9	10-15		
Factor	1 Point	2 Points	3 Points		
Total bilirubin (µmol/L)	<34	34-50	>50		
Serum albumin (g/L)	>35	28-35	<28		
Prothrombin time / international normalized ratio	<1.7	1.71-2.3	>2.3		
Ascites	None	Mild	Moderate to Severe		
Hepatic encephalopathy	None	Grade I-II (or supressed with medication)	Grade III-IV (or refractory)		

Last update: November 6, 2019