



## NS3 Protease Inhibitor + Peginterferon/Ribavirin-Experienced, Genotype 1 Patients With Compensated Cirrhosis

Recommended and alternative regimens listed by evidence level and alphabetically for:

### NS3 Protease Inhibitor (Telaprevir, Boceprevir, or Simeprevir) + Peginterferon/Ribavirin Treatment-Experienced, Genotype 1 Patients With Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	12 weeks	IIa, B
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) plus weight-based ribavirin	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg) plus weight-based ribavirin for all genotype 1b patients, and genotype 1a patients without baseline NS5A RASs <sup>c</sup> for elbasvir	12 weeks	IIa, B
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg) plus weight-based ribavirin for genotype 1a patients with baseline NS5A RASs <sup>c</sup> for elbasvir	16 weeks	IIa, B

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.

<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

<sup>c</sup> Includes genotype 1a RASs at amino acid position 28, 30, 31, or 93 known to confer [antiviral resistance to elbasvir](#). Baseline testing for these RASs is recommended for patients receiving elbasvir/grazoprevir-based regimens.

**Last update:** November 6, 2019