

Testing, Evaluation, and Monitoring of Hepatitis C

The following pages address testing, evaluation, and monitoring of patients with HCV before, during and after antiviral therapy.

- [HCV Testing and Linkage to Care](#)
- [When and in Whom to Initiate HCV Therapy](#)
- [Overview of Cost, Reimbursement, and Cost-Effectiveness Considerations for Hepatitis C Treatment Regimens](#)
- [Monitoring Patients Who Are Starting HCV Treatment, Are on Treatment, or Have Completed Therapy](#)
- [HCV Resistance Primer](#)

Last update: August 27, 2020

HCV Testing and Linkage to Care

One-Time Hepatitis C Testing

Recommendations for One-Time Hepatitis C Testing	
RECOMMENDED	RATING
One-time, routine, opt out HCV testing is recommended for all individuals aged 18 years or older.	I, B
One-time HCV testing should be performed for all persons less than 18 years old with activities, exposures, or conditions or circumstances associated with an increased risk of HCV infection (see below).	I, B
Prenatal HCV testing as part of routine prenatal care is recommended with each pregnancy.	I, B
Periodic repeat HCV testing should be offered to all persons with activities, exposures, or conditions or circumstances associated with an increased risk of HCV exposure (see below).	IIa, C
Annual HCV testing is recommended for all persons who inject drugs , for HIV-infected men who have unprotected sex with men , and men who have sex with men taking pre-exposure prophylaxis (PrEP) .	IIa, C
<p>Risk Activities</p> <ul style="list-style-type: none"> • Injection drug use (current or ever, including those who injected only once) • Intranasal illicit drug use • Men who have sex with men <p>Risk Exposures</p> <ul style="list-style-type: none"> • Persons on long-term hemodialysis (ever) • Persons with percutaneous/parenteral exposures in an unregulated setting • Healthcare, emergency medical, and public safety workers after needlestick, sharps, or mucosal exposure to HCV-infected blood • Children born to HCV-infected women • Recipients of a prior transfusion or organ transplant, including persons who: <ul style="list-style-type: none"> ◦ Were notified that they received blood from a donor who later tested positive for HCV ◦ Received a transfusion of blood or blood components, or underwent an organ transplant before July 1992 ◦ Received clotting factor concentrates produced before 1987 • Persons who were ever incarcerated <p>Other Conditions and Circumstances</p> <ul style="list-style-type: none"> • HIV infection • Sexually active persons about to start pre-exposure prophylaxis (PrEP) for HIV • Chronic liver disease and/or chronic hepatitis, including unexplained elevated alanine aminotransferase (ALT) levels • Solid organ donors (living and deceased) and solid organ transplant recipients 	

Based on the 2013-2016 NHANES data among the general noninstitutionalized US population, an estimated 4.1 million people have had HCV exposure (HCV-antibody-positive), including 2.4 million with active HCV infection (HCV-RNA-positive) ([Hofmeister, 2019](#)). Total HCV burden in the US also includes those not accounted for in NHANES data—incarcerated, institutionalized, or unsheltered homeless persons—with estimates ranging from 380,000 to 800,000 additional HCV-antibody-positive persons ([Hofmeister, 2019](#)); ([Edlin, 2015](#)). Approximately 50% of all infected persons are unaware that they have HCV ([Holmberg, 2013](#)); ([Denniston, 2012](#)).

HCV screening is recommended because of the known benefits of care and treatment in reducing the risk of hepatocellular carcinoma and all-cause mortality, and the potential public health benefit of reducing transmission through early treatment, viral clearance, and reduced risk behaviors ([Chou, 2020](#)); ([Owens, 2020](#)); ([Schillie, 2020](#)); ([Smith, 2012](#)).

HCV is primarily transmitted through percutaneous exposure to infected blood. Other modes of transmission include mother-to-infant and contaminated devices shared for noninjection drug use. Sexual transmission also occurs but is generally inefficient except among HIV-infected men who have unprotected sex with men ([Pakianathan, 2018](#)).

Injection drug use (IDU) poses the greatest risk for HCV infection, accounting for at least 60% of acute HCV infections in the United States. Healthcare exposures are important sources of transmission, including: the receipt of blood products in the US prior to 1992 (after which routine screening of the blood supply was implemented); receipt of clotting factor concentrates in the US before 1987; receipt of blood or blood products in other countries (risk depends on country and screening practices); long-term hemodialysis; needlestick injuries among healthcare workers; and patient-to-patient transmission resulting from poor infection control practices. Other risk factors include having been born to an HCV-infected mother, incarceration, and percutaneous or parenteral exposures in an unregulated setting. Examples include tattoos received outside of licensed parlors and medical procedures performed internationally or domestically where strict infection control procedures may not have been followed (eg, surgery before implementation of universal precautions) ([Hellard, 2004](#)).

The importance of these risk factors might differ based on geographic location and population ([Schillie, 2020](#)); ([Owens, 2020](#)). An estimated 12% to 39% of incarcerated persons in North America are HCV-antibody-positive, supporting the recommendation to test this population for HCV ([Larney, 2013](#)); ([Allen, 2003](#)); ([Weinbaum, 2003](#)).

Because of shared transmission modes, persons with HIV infection are at risk for HCV. Annual HCV testing is recommended for sexually active HIV-infected adolescent and adult men who have sex with men. The presence of concomitant ulcerative sexually transmitted infections, proctitis related to sexually transmitted infections, or high-risk sexual or drug use practices may warrant more frequent testing. Sexual transmission is particularly a risk for HIV-infected [men who have unprotected sex with men](#) ([Hosein, 2013](#)); ([van de Laar, 2010](#)). Testing sexually active, non-HIV-infected persons for HCV infection before starting and while receiving pre-exposure prophylaxis (PrEP) for HIV prevention should also be considered ([Hoorneborg, 2020](#)); ([Volk, 2015](#)).

Data also support testing in all deceased and living solid organ donors because of the risk of HCV infection posed to the recipient ([Lai, 2013](#)); ([Seem, 2013](#)). Although hepatitis C testing guidelines from the US CDC and the USPSTF do not specifically recommend testing immigrants from countries with a high HCV prevalence (eg, Egypt and Pakistan), such persons 18 years or older are included in the one-time, opt out HCV testing recommendation.

CDC established risk-based HCV testing guidelines in 1998 ([CDC, 1998](#)). These guidelines were expanded in 2012 with a recommendation to offer one-time HCV testing to all persons born from 1945 through 1965 without prior ascertainment of HCV risk factors. This recommendation was supported by evidence demonstrating that a risk-based strategy alone failed to identify >50% of HCV infections, due in part to patient underreporting of their risk and provider limitations in ascertaining risk factor information. The USPSTF also recommended a one-time HCV test in asymptomatic persons belonging to the 1945 through 1965 birth cohort as well as other individuals based on exposures, behaviors, and conditions or circumstances that increase HCV infection risk.

Since the birth cohort recommendation was adopted, however, there has been an increase in the number of acute and chronic HCV infections reported in individuals born after 1965 ([Zibbell, 2018](#)); ([Ly, 2017](#)); ([Suryaprasad, 2014](#)). The increase in HCV incidence and prevalence among a younger cohort results from the opioid epidemic and increased IDU.

This shift in HCV epidemiology and the known failures of risk-based testing warranted an expansion of the recommendation. Accordingly, CDC updated screening recommendations in 2020 to include HCV screening at least once in a lifetime for all adults aged ≥ 18 years as well as HCV screening of all pregnant women during each [pregnancy](#), except in settings where the prevalence of HCV infection (ie, HCV-RNA positivity) is $< 0.1\%$. Both recommendations were based on extensive literature review and estimates of cost-effectiveness of screening. In their recommendation, CDC noted that no states have an estimated HCV-RNA prevalence $< 0.1\%$ ([Schillie, 2020](#)). USPSTF also recently issued recommendations for one-time, routine, opt out testing of adults aged 18 through 79 years ([Owens, 2020](#)).

For the CDC 2020 testing guidelines, a systematic review included a harms assessment for hepatitis C screening during pregnancy. This review was augmented by input from subject matter experts, studies not captured through the formal literature review, and the peer-review process. Despite several plausible harms (including insurability and employability issues, legal ramifications and potential loss of infant custody, unnecessary cesarean deliveries, and unnecessary avoidance of breastfeeding), CDC concluded that identified or potential harms did not outweigh the benefits of HCV screening ([Schillie, 2020](#)).

Generally, routine HCV testing is cost-effective because of increasing HCV incidence and prevalence among people who inject drugs (PWID) and the decreasing cost of DAA therapy. Many patients at greatest risk for HCV infection and transmission do not readily report their highly stigmatized risk activities. Studies conducted in US urban emergency departments, for example, reveal that 15% to 25% of patients with previously unidentified HCV infection were born after 1965 and/or have no reported history of IDU and are, therefore, missed by even perfect implementation of risk-based testing guidance ([Schechter-Perkins, 2018](#)); ([Hsieh, 2016](#)); ([Lyons, 2016](#)). Reinfection among those actively using drugs is common, but because HCV testing is a low-cost intervention and therapy is both highly effective and cost-effective, routine testing provides good economic value (ie, cost-effectiveness) even when many people need to be tested and treated more than once during their lifetime.

Several cost-effectiveness studies published since release of the birth cohort recommendations have demonstrated that routine, one-time HCV testing among all adults in the US would likely identify a substantial number of HCV cases that are currently being missed, and that doing so would be cost-effective. One research group employed simulation modeling to compare several versions of routine guidance, including routine testing for adults aged ≥ 40 years, ≥ 30 years, and ≥ 18 years. The investigators found that routine HCV testing for all adults ≥ 18 years was cost-effective compared to current guidance, and potentially cost-saving compared to testing only those aged ≥ 30 years or ≥ 40 years ([Barocas, 2018](#)). The study further demonstrated that routine testing remained cost-effective unless HCV infection had no impact on healthcare utilization and no impact on quality of life. Another research team similarly found that routine HCV testing for all adults aged ≥ 18 years is likely cost-effective compared to current guidance, provided the HCV prevalence among those born after 1965 is $> 0.07\%$ ([Eckman, 2019](#)). Notably, these studies reached similar conclusions despite being conducted independently and employing different simulation modeling approaches. Further, a variety of studies have tested the cost-effectiveness of routine HCV testing in specific venues, including correctional settings ([He, 2016](#)), substance use treatment centers ([Schackman, 2018](#)); ([Schackman, 2015](#)), and federally qualified health centers ([Assoumou, 2018](#)). All of these studies demonstrated that routine HCV testing and treatment was cost-effective, even when linkage to HCV treatment after testing was poor and the rate of HCV reinfection among injection drug users was high.

Analyses focusing on pregnant women have demonstrated similar findings. One analysis calculated an incremental cost-effectiveness ratio (ICER) of \$2,826 per quality-adjusted life-year (QALY) gained for universal screening of pregnant women compared with risk-based screening at an HCV-RNA positivity prevalence of 0.73% ([Chaillon, 2019](#)). Although real-world data informing screening during each pregnancy are lacking, a modeled analysis suggested that hepatitis C screening during each pregnancy would be cost-effective. Using a hepatitis C prevalence of 0.38% among pregnant women, the analysis found that universal hepatitis C screening during the first trimester of each pregnancy compared with the current practice of risk-based screening had an ICER of \$41,000 per QALY gained ([Tasillo, 2019](#)). Universal screening reduced HCV-attributable mortality by 16% and increased the proportion of infants identified as HCV-exposed, from 44% to 92%. ICER remained at or below \$100,000 per QALY gained if hepatitis C prevalence was higher than 0.16% ([Schillie, 2020](#)).

Evidence regarding the frequency of HCV testing in persons at risk for ongoing exposures to the virus is lacking. Clinicians should, therefore, determine the periodicity of testing based on the risk of infection or reinfection. Because of the high incidence of HCV infection among [PWID](#) and HIV-infected [men who have unprotected sex with men](#), HCV testing at least

annually using an assay that detects HCV RNA (ie, a quantitative HCV-RNA test) if they have been previously exposed, is recommended among such individuals ([Newsom, 2017](#)); ([Aberg, 2014](#)); ([Witt, 2013](#)); ([Bravo, 2012](#)); ([Linac, 2012](#)); ([Wandeler, 2012](#)); ([Williams, 2011](#)).

Implementation of clinical decision support tools or prompts for HCV testing in [electronic health records](#) could facilitate reminding clinicians of HCV testing when indicated ([Hsu, 2013](#)); ([Litwin, 2012](#)).

Initial HCV Testing and Follow-Up

Recommendations for Initial HCV Testing and Follow-Up	
RECOMMENDED	RATING
HCV-antibody testing with reflex HCV RNA polymerase chain reaction (PCR) testing is recommended for initial HCV testing.	I, A
Among persons with a negative HCV-antibody test who were exposed to HCV within the prior 6 months, HCV-RNA or follow-up HCV-antibody testing 6 months or longer after exposure is recommended. HCV-RNA testing can also be considered for immunocompromised persons.	I, C
Among persons at risk of reinfection after previous spontaneous or treatment-related viral clearance, HCV-RNA testing is recommended because a positive HCV-antibody test is expected.	I, C
Quantitative HCV-RNA testing is recommended prior to initiation of antiviral therapy to document the baseline level of viremia (ie, baseline viral load).	I, A
HCV genotype testing may be considered for those in whom it may alter treatment recommendations.	I, A
Persons found to have a positive HCV-antibody test and negative results for HCV RNA by PCR should be informed that they do not have evidence of current (active) HCV infection but are not protected from reinfection.	I, A

All persons for whom HCV screening is recommended should initially be tested for HCV antibody ([CDC, 2013](#)); ([Alter, 2003](#)) using an assay approved by the US Food and Drug Administration (FDA). FDA-approved tests include laboratory-based assays and a point-of-care assay (ie, OraQuick™ HCV Rapid Antibody Test [OraSure Technologies]) ([Lee, 2011](#)). The latter is an indirect immunoassay with a sensitivity and specificity similar to those of laboratory-based HCV-antibody assays. Point-of-care assays are valuable in the community setting and allow for sample collection with a finger stick rather than standard phlebotomy. If point-of-care assays are used, reporting of results to the medical record and health authorities should follow protocols used for laboratory-based HCV-antibody tests. When possible, positive point-of-care antibody tests should be followed-up with immediate HCV-RNA confirmatory testing rather than referring the patient to another provider or setting to have the test performed. Table 1 lists FDA-approved, commercially available HCV-antibody screening assays.

Table 1. FDA-Approved HCV-Antibody Screening Assays

Assay	Manufacturer	Format
Abbott HCV EIA 2.0	Abbott Laboratories Abbott Park, IL, USA	EIA ^a (manual)

Assay	Manufacturer	Format
Advia Centaur™ HCV Assay	Siemens Healthcare Malvern, PA, USA	CLIA ^b (automated)
Architect Anti-HCV	Abbott Laboratories Abbott Park, IL, USA	CMIA ^c (automated)
AxSYM™ Anti-HCV	Abbott Laboratories Abbott Park, IL, USA	MEIA ^d (automated)
Elecsys™ Anti-HCV II	Roche Diagnostics Indianapolis, IN, USA	ECLIA ^e (automated)
OraQuick™ HCV Rapid Antibody Test	OraSure Technologies, Inc. Bethlehem, PA, USA	Immunochromatographic (manual)
Ortho HCV Version 3.0 ELISA Test System	Ortho-Clinical Diagnostics, Inc. Raritan, NJ, USA	EIA ^a (manual)
Vitros Anti-HCV	Ortho-Clinical Diagnostics, Inc. Rochester, NY, USA	CLIA ^b (automated)
^a EIA: enzyme immunoassay ^b CLIA: chemiluminescent immunoassay ^c CMIA: chemiluminescent microparticle immunoassay ^d MEIA: microparticle enzyme immunoassay ^e ECLIA: electrochemiluminescent immunoassay		


A positive HCV-antibody test indicates current (active) HCV infection (acute or chronic), past infection that has resolved, or rarely a false positive result ([Pawlotsky, 2002](#)). A test to detect HCV viremia is therefore necessary to confirm active HCV infection and guide clinical management, including initiation of HCV treatment. Many reference laboratories now offer HCV-antibody testing that automatically reflexes to HCV-RNA PCR testing if the antibody test is positive. This should be considered the optimal testing approach in a clinical setting because it requires only a single blood draw without the need to bring people back to care for confirmatory testing, a major barrier in the continuum of care ([Mera, 2016](#)). Collection of dried blood spot (DBS) samples also allows for assessment of HCV antibodies and reflex HCV-RNA testing by testing spots sequentially. DBS samples can be collected using a finger stick rather than phlebotomy and can be transported without an intact cold chain, making it useful in rural areas and in people for whom phlebotomy may be a testing barrier ([Lange, 2017](#)).

HCV-RNA testing should also be performed in persons with a negative HCV-antibody test who are either immunocompromised (eg, persons receiving chronic hemodialysis) ([KDIGO, 2008](#)) or might have been exposed to HCV within the last 6 months because these persons may be HCV antibody negative. An HCV-RNA test is also needed to detect reinfection in HCV-antibody-positive persons after previous spontaneous or treatment-related viral clearance.

Detection of HCV core antigen in the blood also indicates active HCV infection. Because the sensitivity of HCV core antigen testing is less than that of HCV-RNA testing, if an HCV core antigen test is used to assess viremia, antibody-positive samples that test negative for HCV core antigen should have a confirmatory HCV-RNA test to exclude a false negative core antigen result ([van Tilborg, 2018](#)).

An FDA-approved quantitative or qualitative HCV-RNA test with a detection level of ≤ 25 IU/mL should be used to detect HCV RNA. Figure 1 shows the CDC-recommended HCV testing algorithm.

Figure 1. CDC-Recommended Testing Sequence for Identifying Current HCV Infection



^a For persons who might have been exposed to HCV within the past 6 months, testing for HCV RNA or follow-up testing for HCV antibody should be performed. For persons who are immunocompromised, testing for HCV RNA should be performed.

^b To differentiate past, resolved HCV infection from biologic false positivity for HCV antibody, testing with another HCV-antibody assay can be considered. Repeat HCV-RNA testing if the person tested is suspected to have had HCV exposure within the past 6 months or has clinical evidence of HCV disease, or if there is concern regarding the handling or storage of the test specimen.

Adapted from Centers for Disease Control and Prevention ([CDC, 2013](#)).


Persons who have a positive HCV-antibody test and negative results for HCV RNA by PCR should be informed that they do not have laboratory evidence of current HCV infection. Additional HCV testing is typically unnecessary. The HCV-RNA test can be repeated when there is a high index of suspicion for recent infection or in patients with ongoing HCV infection risk.

Clinicians (or patients) may seek additional testing to determine whether a positive HCV-antibody test represents a remote, resolved HCV infection or a false positive. For patients with no apparent risk for HCV infection, the likelihood of a false positive HCV-antibody test is directly related to the HCV prevalence in the tested population. False positive HCV-antibody tests most commonly occur in populations with a low prevalence of HCV infection ([Alter, 2003](#)). If further testing is desired to distinguish between a true positive vs biologic false positivity for HCV antibody, repeat testing may be done with a different FDA-approved, HCV-antibody assay. A biologic false result should not occur with 2 different assays because they target different regions of the virus, making it very unlikely that both would falsely detect a cross-reactive antigen ([CDC, 2013](#)); ([Vermeersch, 2008](#)).

Prior to initiation of antiviral therapy, quantitative HCV-RNA testing should be used to determine the baseline level of viremia (ie, viral load), which may affect treatment duration with certain regimens. The degree of viral load decline after initiation of treatment is less predictive of sustained virologic response (SVR) in the era of direct-acting antiviral (DAA) therapy compared to previous interferon-based treatment (see [Pretreatment and On-Treatment Monitoring](#)).

With the advent of pangenotypic HCV treatment regimens, HCV genotyping is no longer required prior to treatment initiation for all individuals. In those with evidence of cirrhosis and/or past unsuccessful HCV treatment, treatment regimens may differ by genotype and thus pretreatment genotyping is recommended (see [Treatment-Naive](#) and [Treatment-Experienced](#) sections). For noncirrhotic treatment-naive patients, although genotyping may impact the preferred treatment approach, it is not required if a pangenotypic regimen is used (see [Simplified Treatment Algorithm](#)).

Counseling Persons With Active HCV Infection

Recommendations for Counseling Persons With Active HCV Infection	
RECOMMENDED	RATING 
Persons with current HCV infection should receive education and interventions aimed at reducing liver disease progression and preventing HCV transmission.	IIa, B
Abstinence from alcohol and, when appropriate, interventions to facilitate cessation of alcohol consumption should be advised for all persons with HCV infection.	IIa, B
Evaluation for other conditions that may accelerate liver fibrosis, including hepatitis B and HIV infections, is recommended for all persons with active HCV infection.	IIb, B
Evaluation for advanced fibrosis using noninvasive tests (serum panels, elastography) or liver biopsy, if required, is recommended for all persons with HCV infection to facilitate an appropriate decision regarding HCV treatment strategy, and to determine the need for initiating additional measures for cirrhosis management (eg, hepatocellular carcinoma screening) (see Monitoring section).	I, A
Vaccination against hepatitis A and hepatitis B is recommended for all susceptible persons with HCV infection.	IIa, C
Vaccination against pneumococcal infection is recommended for all patients with cirrhosis.	IIa, C
All persons with HCV infection should be provided education about how to prevent HCV transmission	I, C

Recommendations for Counseling Persons With Active HCV Infection

to others.

In addition to receiving antiviral therapy, HCV-infected persons should be educated about how to prevent further liver damage. Most important is prevention of the potential deleterious effect of alcohol. Numerous studies have found a strong association between excess alcohol use and the development or progression of liver fibrosis, and the development of hepatocellular carcinoma ([Safdar, 2004](#)); ([Harris, 2001](#)); ([Bellentani, 1999](#)); ([Corrao, 1998](#)); ([Wiley, 1998](#)); ([Poynard, 1997](#)); ([Noda, 1996](#)).

Daily consumption of >50 g of alcohol has a high likelihood of worsening fibrosis. Some studies indicate that daily consumption of lesser amounts of alcohol also exerts a deleterious effect on the liver; these data, however, are controversial ([Hagström, 2017](#)); ([Younossi, 2013b](#)); ([Westin, 2002](#)). Persons who abuse alcohol and have alcohol dependence require treatment and consideration for referral to an addiction specialist.

Hepatitis B virus (HBV) and HIV coinfection have been associated with a poorer HCV prognosis in cohort studies ([Puoti, 2017b](#)); ([Kruse, 2014](#)); ([Thein, 2008a](#)); ([Zarski, 1998](#)). Because of overlapping risk factors for these infections and benefits associated with their identification and treatment, HCV-infected persons should be tested for HIV antibody and hepatitis B surface antigen (HBsAg) using standard screening assays ([Moyer, 2013](#)); ([CDC, 2008](#)). (See [USPSTF HIV screening recommendations](#) and [CDC hepatitis B screening recommendations](#).) Persons who test positive for HBsAg require monitoring during HCV treatment because of HBV reactivation risk ([Lee, 2018](#)). Anti-HBV therapy may also be considered (see [reactivation of HBV in the Monitoring section](#)). For persons who test negative for HBsAg but positive for hepatitis B core antibodies (anti-HBc), with or without hepatitis B surface antibodies (anti-HBs), have resolved HBV infection; the risk of clinically significant HBV reactivation with HCV therapy is very low and no further workup is required ([Mücke, 2018](#)). Patients should be counseled about how to reduce their risk of acquiring these infections and HBV vaccination is recommended when appropriate.

Assessment of Liver Disease Severity

The severity of liver disease associated with chronic HCV infection is a key factor in determining the initial and follow-up evaluation of patients. Noninvasive tests using serum biomarkers, elastography, or liver imaging allow for accurate diagnosis of cirrhosis in most individuals (see [pretreatment workup in When and in Whom to Initiate HCV Therapy](#)). Liver biopsy is rarely required but may be considered if other causes of liver disease are suspected.

Noninvasive methods frequently used to estimate liver disease severity include:

- Liver-directed physical exam (normal in most patients)
- Routine blood tests (eg, ALT, AST, albumin, bilirubin, international normalized ratio [INR], and CBC with platelet count)
- Serum fibrosis marker panels
- Transient elastography
- Liver imaging (eg, ultrasound or CT scan)

Simple calculations derived from routine blood tests—such as the serum AST-to-platelet ratio index (APRI) ([Wai, 2003](#)) and FIB-4 score ([Sterling, 2006](#))—as well as assessment of liver surface nodularity and spleen size by liver ultrasound or other cross-sectional imaging modalities can help determine if patients with HCV have cirrhosis and associated portal hypertension. The presence of portal hypertension is associated with a greater likelihood of developing future hepatic complications in untreated patients ([Chou, 2013](#)); ([Rockey, 2006](#)). Elastography provides instant information regarding liver stiffness and vibration-controlled transient elastography at the point of care and can reliably distinguish patients with a high vs low likelihood of cirrhosis ([Bonder, 2014](#)); ([Castera, 2012](#)). A more detailed discussion regarding fibrosis

assessment is found in the [When and In Whom to Initiate Therapy](#) section.


Persons with known or suspected bridging fibrosis and cirrhosis are at increased risk for developing complications of advanced liver disease and require frequent follow-up. They should also avoid hepatotoxic drugs, such as excessive acetaminophen (>2 g/d) and certain herbal supplements. Nephrotoxic drugs, such as nonsteroidal anti-inflammatory drugs, should also be avoided. Ongoing imaging surveillance for liver cancer and gastroesophageal varices is also recommended for these patients ([Fontana, 2010](#)); ([Sangiovanni, 2006](#)). Persons with cirrhosis are more susceptible to invasive pneumococcal infection ([Marrie, 2011](#)) and should receive pneumococcal vaccination ([CDC, 2012](#)).

Exposure to infected blood is the primary mode of HCV transmission. HCV-infected persons must be informed of the precautions needed to avoid exposing others to infected blood. This is particularly important for PWID given that HCV transmission in this population primarily results from sharing needles and other contaminated drug injection equipment. Epidemics of acute HCV due to sexual transmission in HIV-infected men who have sex with men have also been described ([Urbanus, 2009](#)); ([van de Laar, 2009](#)); ([Fierer, 2008](#)). Table 2 outlines measures to avoid HCV transmission. HCV is not spread by sneezing, hugging, holding hands, coughing, or sharing eating utensils or drinking glasses, nor is it transmitted through food or water.

Table 2. Measures to Prevent HCV Transmission

HCV-infected persons should be counseled to avoid sharing toothbrushes and dental or shaving equipment, and be cautioned to cover any bleeding wound to prevent the possibility of others coming into contact with their blood.
Persons should be counseled about harm reduction related to illicit drug use, including entering substance use treatment programs. Those who continue to inject drugs should be counseled to: <ul style="list-style-type: none"> • Avoid reusing or sharing syringes, needles, water, cotton, and other drug preparation equipment. • Use new sterile syringes and filters, and disinfected cookers. • Clean the injection site with a new alcohol swab. • Dispose of syringes and needles after 1 use in a safe, puncture-proof container.
Persons with HCV infection should be advised not to donate blood and to discuss HCV serostatus prior to donation of body organs, other tissue, or semen.
Persons with HIV infection and those with multiple sexual partners or sexually transmitted infections should be encouraged to use barrier precautions to prevent sexual transmission. Other persons with HCV infection should be counseled that the risk of sexual transmission is low and may not warrant barrier protection.
Household surfaces and implements contaminated with visible blood from an HCV-infected person should be cleaned using a dilution of 1 part household bleach to 9 parts water. Gloves should be worn when cleaning up blood spills.

Linkage to Care

Recommendation for Linkage to Care	
RECOMMENDED	RATING 
All persons with active HCV infection should be linked to a healthcare provider who is knowledgeable in and prepared to provide comprehensive management.	Ila, C

Improved identification of active HCV infection and treatment advances will have limited impact on HCV-related morbidity and mortality without concomitant improvement in linkage to care. All patients with current HCV infection and a positive HCV-RNA test should be evaluated by a healthcare provider with expertise in assessment of liver disease severity and HCV treatment. Subspecialty care and consultation may be required for persons with HCV infection who have advanced fibrosis or cirrhosis (Metavir stage \geq F3), including possible referral for consideration of liver transplantation in those with evidence of hepatic decompensation.

Only an estimated 13% to 18% of HCV-infected persons in the US had received treatment by 2013 ([Holmberg, 2013](#)). Lack of appropriate clinician assessment and delays in linkage to care can result in negative health outcomes. Furthermore, patients who are lost to follow-up fail to benefit from evolving evaluation and treatment options.

Commonly cited patient-related barriers to treatment initiation include contraindications to treatment (eg, medical or psychiatric comorbidities); lack of acceptance of treatment (eg, asymptomatic nature of disease, competing priorities, low treatment efficacy, long treatment duration, and adverse effects); and lack of access to treatment (eg, cost and distance to specialist) ([Clark, 2012](#)); ([Arora, 2011](#)); ([Khokhar, 2007](#)).

Common healthcare provider-related barriers include perceived patient-related barriers (eg, fear of adverse effects, treatment duration, cost, and effectiveness); lack of expertise in HCV treatment; lack of specialty referral resources; resistance to treating persons currently using illicit drugs or alcohol; and concern about HCV treatment cost ([Reilley, 2014](#)); ([McGowan, 2013](#)); ([Morrill, 2005](#)).

Data do not support exclusion of HCV-infected persons from consideration for hepatitis C therapy based alcohol intake or use of illicit drugs (see [Identification and Management of HCV in People Who Inject Drugs](#)). Some possible strategies to address HCV treatment barriers are listed in Table 3.

Table 3. Common Barriers to and Misconceptions Regarding HCV Treatment and Potential Strategies

Barrier	Strategy
Comorbid conditions (eg, substance use psychiatric disorders, uncontrolled chronic medical conditions)	<ul style="list-style-type: none"> • Conduct counseling and education. • Refer for services (eg, mental health services, medications for opioid use disorder [MOUDs], and syringe service programs). • Co-localize services (eg, primary care, medical homes, and drug treatment).
Competing priorities and loss to follow-up	<ul style="list-style-type: none"> • Conduct counseling and education. • Engage case managers and patient navigators. Consider other strategies such as incentives, peer navigators, and transportation assistance. • Co-localize services (eg, primary care, medical homes, and drug treatment).
Long treatment duration and adverse effects	<ul style="list-style-type: none"> • Conduct counseling and education. • Consider other strategies like incentives, peer navigators, and transportation assistance. • Utilize directly observed therapy.
Lack of access to treatment (eg, out-of-pocket costs, high copays, lack of insurance, geographic distance, and/or lack of specialist availability)	<ul style="list-style-type: none"> • Leverage expansion of coverage through the Patient Protection and Affordable Care Act. • Participate in models of care involving close collaboration between primary care clinicians and specialists. • Liaise with pharmaceutical patient assistance programs and copay assistance programs. • Co-localize services (eg, primary care, medical homes, and drug treatment).
Lack of practitioner expertise	<ul style="list-style-type: none"> • Collaborate with specialists (eg, project ECHO-like models and telemedicine). • Develop accessible, clear HCV treatment guidelines. • Develop electronic health record performance measures and clinical decision support tools (eg, pop-up reminders and standing orders).

Co-localization of HCV screening, evaluation, and treatment with other medical or social services (ie, integrated care) is a strategy that addresses several treatment barriers. Co-localization has already been applied to settings with high HCV prevalence (eg, correctional facilities, needle exchange programs, substance abuse treatment centers, and harm reduction programs) but this type of care is not uniformly available ([Burton, 2019](#)); ([Harrison, 2019](#)); ([Morey, 2019](#)); ([Schulkind, 2019](#)); ([Bruggmann, 2013](#)); ([Islam, 2012](#)); ([Stein, 2012](#)). A recent study demonstrated that integrated care—consisting of multidisciplinary care coordination and patient case management—increased the proportion of patients with HCV infection and psychiatric illness or substance use who begin antiviral therapy and achieve SVR without serious adverse events ([Ho, 2015](#)).

A strategy that addresses lack of access to specialists—a primary barrier to hepatitis C care—is participation in models

involving close collaboration between primary care practitioners and subspecialists ([Beste, 2017b](#)); ([Rossaro, 2013](#)); ([Miller, 2012](#)); ([Arora, 2011](#)). Such collaborations have used telemedicine and knowledge networks to overcome geographic distances to specialists ([Rossaro, 2013](#)); ([Arora, 2011](#)) or the availability of experienced providers in a methadone or correctional setting ([Morey, 2019](#)); ([Talal, 2019](#)). For example, project ECHO ([Extension for Community Healthcare Outcomes](#)) uses videoconferencing to enhance primary care practitioner capacity in rendering HCV care and treatment to New Mexico's large rural and underserved population ([Arora, 2011](#)). Through case-based learning and real-time feedback from a multidisciplinary team of specialists (gastroenterology, infectious disease, pharmacology, and psychiatry practitioners), project ECHO has expanded HCV treatment access in populations that might have otherwise remained untreated. The short duration of treatment and few serious adverse events associated with DAA therapy present an opportunity to expand the number of primary care providers engaged in HCV management and treatment. This expansion will support the goals of HCV elimination and overcome barriers associated with the need for subspecialty referrals. The ASCEND trial utilized a real-world cohort of patients at urban federally qualified health centers and found that HCV treatment administered by nonspecialist providers was as safe and effective as that provided by specialists ([Kattakuzhy, 2017](#)).

Additional strategies for enhancing linkage to and retention in care could be adapted from other fields, such as tuberculosis and HIV. For example, use of directly observed therapy has enhanced adherence to tuberculosis treatment, and use of case managers and patient navigators has reduced loss of follow-up in HIV care ([Govindasamy, 2012](#)). Recent hepatitis C testing and care programs have identified the use of patient navigators or care coordinators as important interventions in overcoming challenges associated with linkage to and retention in care ([Ford, 2018](#)); ([Coyle, 2015](#)); ([Trooskin, 2015](#)). There are also data suggesting that financial incentives and peer navigation may be useful to support treatment adherence in patients with substance use disorders ([Ward, 2019](#)); ([Wohl, 2017](#)). Ongoing assessment of efficacy and comparative effectiveness of this and additional strategies is a crucial area of future research for patients with HCV infection. Replication and expansion of best practices and new models for linkage to HCV care will also be crucial to maximize the public health impact of newer treatment paradigms.

Last update: August 27, 2020

When and in Whom to Initiate HCV Therapy


Successful hepatitis C treatment results in sustained virologic response (SVR), which is tantamount to virologic cure and, as such, is expected to benefit nearly all chronically infected persons. When the US Food and Drug Administration (FDA) approved the first interferon-sparing treatment for HCV infection, many patients who had previously been “warehoused” sought treatment. The infrastructure (ie, experienced practitioners, budgeted healthcare dollars, etc) did not yet exist to treat all patients immediately. Thus, the panel offered guidance for prioritizing treatment first for those with the greatest need.

Since that time, there have been opportunities to treat many of the highest-risk patients and accumulate real-world experience regarding the tolerability and safety of interferon-free HCV regimens. More importantly, from a medical standpoint, data continue to accumulate that demonstrate the many benefits, both intrahepatic and extrahepatic, that accompany HCV eradication. Therefore, the panel continues to recommend treatment for all patients with chronic HCV infection, except those with a short life expectancy that cannot be remediated by HCV treatment, liver transplantation, or another directed therapy. Accordingly, prioritization tables have been removed from this section.


Despite the strong recommendation for treatment of nearly all HCV-infected patients, pretreatment assessment of a patient's understanding of treatment goals and provision of education about adherence and follow-up are essential. A well-established therapeutic relationship between clinician and patient remains crucial for optimal outcomes with direct-acting antiviral (DAA) therapies. Additionally, in certain settings there remain factors that impact access to medications and the ability to deliver them to patients. The descriptions of unique populations discussed in this section may help physicians make more informed treatment decisions for these groups. For additional information, see unique patient populations:

[Patients With HIV/HCV Coinfection](#), [Patients With Decompensated Cirrhosis](#), [Patients Who Develop Recurrent HCV Infection Post Liver Transplantation](#), [Patients With Renal Impairment](#), [HCV During Pregnancy](#) and in [Children](#), [Acute HCV Infection](#), and [HCV Post Kidney Transplant](#).

Goal of Treatment

RECOMMENDED	RATING 
The goal of treatment of HCV-infected persons is to reduce all-cause mortality and liver-related health adverse consequences, including end-stage liver disease and hepatocellular carcinoma, by the achievement of virologic cure as evidenced by a sustained virologic response.	I, A

Recommendation for When and in Whom to Initiate Treatment

RECOMMENDED	RATING 
Treatment is recommended for all patients with acute or chronic HCV infection, except those with a short life expectancy that cannot be remediated by HCV therapy, liver transplantation, or another directed therapy. Patients with a short life expectancy owing to liver disease should be managed in consultation with an expert.	I, A

Clinical Benefit of Cure

The proximate goal of HCV therapy is SVR (virologic cure), defined as the continued absence of detectable HCV RNA for at least 12 weeks after completion of therapy. SVR is a marker for cure of HCV infection and has been shown to be durable in large prospective studies in more than 99% of patients followed-up for ≥ 5 years ([Swain, 2010](#)); ([Manns, 2013](#)). While follow-up studies after cure using DAAs are limited, durability of SVR appears to be just as high ([Sarrazin, 2017](#)); ([Reddy, 2018](#)). Patients in whom SVR is achieved have HCV antibodies but no longer have detectable HCV RNA in serum, liver tissue, or mononuclear cells, and achieve substantial improvement in liver histology ([Marcellin, 1997](#)); ([Coppola, 2013](#)); ([Garcia-Bengochea, 1999](#)). Assessment of viral response, including documentation of SVR, requires use of an FDA-approved quantitative or qualitative nucleic acid test (NAT) with a detection level of ≤ 25 IU/mL.

Patients who are cured of their HCV infection experience numerous health benefits, including a decrease in liver inflammation as reflected by improved aminotransferase levels (ie, alanine aminotransferase [ALT] and aspartate aminotransferase [AST]), and a reduction in the rate of liver fibrosis progression ([Poynard, 2002b](#)). Among 3,010 treatment-naïve patients from 4 randomized trials who had pretreatment and post-treatment liver biopsies (separated by a mean of 20 months) and were treated with 10 different interferon-based regimens, 39% to 73% of participants who achieved SVR had improvement in liver fibrosis and necrosis ([Poynard, 2002b](#)). Additionally, cirrhosis resolved in 49% of the cases. Portal hypertension, splenomegaly, and other clinical manifestations of advanced liver disease also improved. Among HCV-infected persons, SVR is associated with a $>70\%$ reduction in the risk of liver cancer (hepatocellular carcinoma [HCC]), and a 90% reduction in the risk of liver-related mortality and liver transplantation ([Morgan, 2013](#)); ([van der Meer, 2012](#)); ([Veldt, 2007](#)).

Cure of HCV infection also reduces symptoms and mortality from severe extrahepatic manifestations, including cryoglobulinemic vasculitis, a condition affecting 10% to 15% of HCV-infected patients ([Fabrizi, 2013](#)); ([Landau, 2010](#)); ([Sise, 2016](#)). HCV-infected persons with non-Hodgkin lymphoma and other lymphoproliferative disorders achieve complete or partial remission in up to 75% of cases following successful therapy for HCV infection ([Gisbert, 2005](#)); ([Takahashi, 2012](#)); ([Svoboda, 2005](#)); ([Mazzaro, 2002](#)); ([Hermine, 2002](#)). These reductions in disease severity contribute to dramatic reductions in all-cause mortality ([van der Meer, 2012](#)); ([Backus, 2011](#)). Furthermore, patients who achieve

SVR have a substantially improved quality of life, which spans their physical, emotional, and social health ([Boscarino, 2015](#)); ([Neary, 1999](#)); ([Younossi, 2014b](#)); ([Gerber, 2016](#)). Conversely, patients who do not achieve SVR after treatment have a continued worsening in health-related quality of life ([Younossi, 2019](#)).

Despite convincing data from observational studies demonstrating the benefit of SVR on all-cause and liver-related mortality, the lack of randomized, placebo-controlled trials of HCV DAA treatment focusing on clinical endpoints (eg, mortality, HCC, liver decompensation, etc) and reliance on surrogate endpoints (eg, HCV RNA) have led some to question the benefits of HCV treatment. In further support of the dramatic benefit of HCV cure, a French cohort study that prospectively followed almost 10,000 patients with chronic HCV infection (including 2,500 who remained untreated for HCV) for a median of 33 months demonstrated a 52% reduction in all-cause mortality and a 34% reduction in HCC ([Carrat, 2019](#)).

Because of the many benefits associated with successful HCV treatment, clinicians should treat HCV-infected patients with antiviral therapy with the goal of achieving SVR, preferably early in the course of chronic hepatitis C before the development of severe liver disease and other complications.

Benefits of Treatment at Early Fibrosis Stages (Metavir Stage Less Than F2)

Initiating therapy in patients with lower-stage fibrosis augments the benefits of SVR. In a long-term follow-up study, 820 patients with biopsy-confirmed Metavir stage F0 or F1 fibrosis were followed for up to 20 years ([Jezequel, 2015](#)). The 15-year survival rate was significantly better for those who experienced SVR than for those whose treatment failed or those who remained untreated (93%, 82%, and 88%, respectively; $P = .003$). The study results argue for consideration of earlier initiation of treatment. Several modeling studies also suggest a greater mortality benefit if treatment is initiated at fibrosis stages prior to F3 ([Øvrehus, 2015](#)); ([Zahnd, 2016](#)); ([Matsuda, 2016](#)).

Treatment delay may decrease the benefit of SVR. In a report from France, 820 patients with biopsy-confirmed Metavir stage F0 or F1 fibrosis were followed for as long as 20 years ([Jezequel, 2015](#)). The authors noted rapid progression of fibrosis in 15% of patients during follow-up, and in patients treated successfully, long-term survival was better. Specifically, at 15 years, survival rate was 92% for those with SVR versus 82% for treatment failures and 88% for those not treated. In a Danish regional registry study, investigators modeled treatment approaches with the aim of evaluating the benefit to the region in terms of reductions in morbidity and mortality and HCV prevalence ([Øvrehus, 2015](#)). Although they note that in their situation of low HCV prevalence (0.4%) with approximately 50% undiagnosed, a policy that restricts treatment to those with Metavir fibrosis stage F3 or higher would decrease mortality from HCC and cirrhosis, the number needed to treat to halve the prevalence of the disease is lower if all eligible patients receive treatment at diagnosis.

A modeling study based on the Swiss HIV cohort study also demonstrated that waiting to treat HCV infection until Metavir fibrosis stages F3 and F4 resulted in 2- and 5-times higher rates of liver-related mortality, respectively, compared with treating at Metavir stage F2 ([Zahnd, 2016](#)). A US Veterans Administration dataset analysis that used very limited endpoints of virologic response dating from the interferon-treatment era suggested that early initiation of therapy (at a fibrosis-4 [FIB-4] score of <3.25) increased the benefit attained with respect to likelihood of treatment success and mortality reduction, and ultimately decreased the number of patients needed to treat to preserve 1 life by almost 50% ([Matsuda, 2016](#)).

Considerations in Specific Populations

Despite the recommendation for treatment of nearly all patients with HCV infection, it remains important for clinicians to understand patient- and disease-related factors that place individuals at risk for HCV-related complications (liver and extrahepatic) as well as for HCV transmission. Although these groups are no longer singled out for high prioritization for treatment, it is nonetheless important that clinicians recognize the unique dimensions of HCV disease and its natural history in these populations. The discussions offered below may assist clinicians in making compelling cases for insurance coverage of treatment when necessary.

Persons With Advanced Liver Disease

For persons with advanced liver disease (Metavir stage F3 or F4), the risk of developing complications of liver disease, such as hepatic decompensation (Child-Turcotte-Pugh [CTP] class B or C [Methods Table 3] ⁱ) or HCC, is substantial and may occur in a relatively short timeframe. A large prospective study of patients with cirrhosis resulting from HCV infection examined the risk of decompensation—including HCC, ascites, jaundice, bleeding, and encephalopathy—and found that the overall annual incidence rate was 3.9% ([Sangiovanni, 2006](#)). The National Institutes of Health (NIH)-sponsored HALT-C study included a group of 220 patients with HCV-related cirrhosis who were observed for approximately 8 years. A primary outcome of death, hepatic decompensation, HCC, or an increase in CTP score ≥ 2 occurred at a rate of 7.5% per year ([Everson, 2006](#)); ([Di Bisceglie, 2008](#)). Patients with a CTP score of ≥ 7 experienced a death rate of 10% per year.

Numerous studies have demonstrated that hepatitis C therapy and the achievement of SVR in this population results in dramatic decreases in hepatic decompensation events, HCC, and liver-related mortality ([Morgan, 2013](#)); ([van der Meer, 2012](#)); ([Backus, 2011](#)); ([Dienstag, 2011](#)); ([Berenguer, 2009](#)); ([Mira, 2013](#)). In the HALT-C study, patients with advanced fibrosis secondary to HCV infection who achieved SVR, compared with patients with similarly advanced liver fibrosis who did not achieve SVR, had a decreased need for liver transplantation (HR, 0.17; 95% CI, 0.06-0.46), decreased development of liver-related morbidity and mortality (HR, 0.15; 95% CI, 0.06-0.38), and decreased HCC (HR, 0.19; 95% CI, 0.04-0.80) ([Dienstag, 2011](#)). Importantly, persons with advanced liver disease also require long-term follow-up and HCC surveillance regardless of treatment outcome (see [Monitoring Patients Who Are Starting Hepatitis C Treatment, Are on Treatment, or Have Completed Therapy](#)).

Given the clinical complexity and need for close monitoring, patients with advanced liver disease that has already decompensated (CTP class B or C [Methods Table 3] ⁱ) should be treated by physicians with experience treating HCV in conjunction with a liver transplantation center, if possible (see [Patients with Decompensated Cirrhosis](#)).

Persons Who Have Undergone Liver Transplantation

In HCV-infected individuals, HCV infection of the liver allograft occurs universally in those with viremia at the time of transplantation. Histologic features of hepatitis develop in about 75% of recipients within the first 6 months following liver transplantation ([Neumann, 2004](#)). By the fifth postoperative year, up to 30% of untreated patients have progressed to cirrhosis ([Neumann, 2004](#)); ([Charlton, 1998](#)). A small proportion of patients (4% to 7%) develop an accelerated course of liver injury (cholestatic hepatitis C, associated with very high levels of viremia) with subsequent rapid allograft failure. Recurrence of HCV infection post transplantation is associated with decreased graft survival for recipients with HCV infection compared to recipients who undergo liver transplantation for other indications ([Forman, 2002](#)).

Effective HCV therapy prior to transplantation resulting in SVR (virologic cure) prevents HCV recurrence post transplantation ([Everson, 2003](#)). In addition, complete HCV viral suppression prior to transplantation prevents recurrent HCV infection of the graft in the majority of cases ([Forns, 2004](#)); ([Everson, 2005](#)). Preliminary data from a study of patients with complications of cirrhosis secondary to HCV infection who were wait-listed for liver transplantation (included patients with MELD scores up to 14 and CTP scores up to 8) found that treatment with sofosbuvir and weight-based ribavirin for up to 48 weeks was well tolerated and associated with an overall SVR of 70% post transplant ([Curry, 2015](#)). Post-transplant SVR was nearly universal among patients who had undetectable HCV RNA for 28 days or longer prior to transplantation.

Treatment of established HCV infection post transplantation also yields substantial improvements in patient and graft survival ([Berenguer, 2008](#)); ([Picciotto, 2007](#)). The availability of effective, interferon-free antiviral therapy has addressed the major hurdles to treating HCV recurrence post transplantation—poor tolerability and efficacy. A multicenter, open-label study evaluated the efficacy of sofosbuvir plus ribavirin to induce virologic suppression in 40 patients after liver transplantation with compensated recurrence of HCV infection. Daily sofosbuvir plus ribavirin for 24 weeks achieved SVR12 in 70% of these patients ([Charlton, 2015](#)). No deaths, graft losses, or episodes of rejection occurred. Six patients had serious adverse events, all of which were considered unrelated to the study treatment. There were no drug interactions reported between sofosbuvir and any of the concomitant immunosuppressive agents. In contrast, treatment with sofosbuvir plus ribavirin, with or without peginterferon, in 64 patients with severe, decompensated cirrhosis resulting from recurrence of HCV infection following liver transplantation was associated with an overall SVR12 of 59% and a

mortality rate of 13% ([Forns, 2015](#)). On an intent-to-treat basis, treatment was associated with clinical improvement in 57% and stable disease in 22% of patients. Given the clinical complexity (including drug-drug interactions and the need for close monitoring), patients with a liver transplant should be treated by physicians with experience in treating this population (see [Patients Who Develop Recurrent HCV Infection Post Liver Transplantation](#)).

Persons at Increased Risk for Rapidly Progressive Fibrosis and Cirrhosis

Fibrosis progression is variable across different patient populations as well as within the same individual over time. Many of the components that determine fibrosis progression and development of cirrhosis in an individual are unknown. However, certain factors, such as coinfection with HIV or the hepatitis B virus (HBV) and prevalent coexistent liver diseases (eg, nonalcoholic steatohepatitis [NASH]), are well recognized contributors to accelerated fibrosis progression (see Table below).

HIV/HCV Coinfection

HIV coinfection accelerates fibrosis progression among HCV-infected persons ([Benhamou, 1999](#)); ([Macias, 2009](#)); ([Konerman, 2014](#)), although control of HIV replication and restoration of the CD4 cell count may mitigate this to some extent but the effect is not completely reversed ([Benhamou, 2001](#)); ([Bräu, 2006](#)); ([Lo Re, 2014](#)). Thus, antiretroviral therapy is not a substitute for HCV treatment. In the largest paired-biopsy study, 282 HIV/HCV-coinfected patients with 435 paired biopsies were prospectively evaluated ([Konerman, 2014](#)). Thirty-four percent of patients showed fibrosis progression of at least 1 Metavir stage at a median of 2.5 years. Importantly, 45% of patients with no fibrosis on initial biopsy had progression. Finally, a more rapid progression to death following decompensation combined with lack of widespread access to liver transplantation and poor outcomes following transplantation highlight the need for HCV treatment in this population regardless of current fibrosis stage (see [Patients with HIV/HCV Coinfection](#)) ([Pineda, 2005](#)); ([Merchante, 2006](#)); ([Terrault, 2012](#)).

HBV/HCV Coinfection

The prevalence of HBV/HCV coinfection is estimated at 1.4% in the United States and 5% to 10% globally ([Tyson, 2013](#)); ([Chu, 2008](#)). Persons with HBV/HCV coinfection and detectable viremia of both viruses are at increased risk for disease progression, decompensated liver disease, and the development of HCC. HBV/HCV-coinfected individuals are susceptible to a process called viral interference wherein one virus may interfere with the replication of the other virus. Thus, when treating one or both viruses with antiviral drugs, periodic retesting of HBV DNA and HCV RNA levels during and after therapy is prudent, particularly if only one of the viruses is being treated at a time. Treatment of HCV infection in such cases utilizes the same genotype-specific regimens as are recommended for HCV mono-infection (see [Initial Treatment of HCV Infection](#)). HBV infection in such cases should be treated as recommended for HBV mono-infection ([Lok, 2009](#)).

Other Coexistent Liver Diseases

Persons with other chronic liver diseases who have coincident chronic HCV infection should be considered for HCV therapy given the potential for rapid liver disease progression. An interferon-free regimen is preferred for immune-mediated liver diseases, such as autoimmune hepatitis, because of the potential for interferon-related exacerbation.

Persons With Extrahepatic Manifestations of Chronic HCV Infection

Cryoglobulinemia

Chronic hepatitis C is associated with a syndrome of cryoglobulinemia, an immune complex and lymphoproliferative disorder that leads to arthralgia, fatigue, palpable purpura, renal disease (eg, membranoproliferative glomerulonephritis), neurologic disease (eg, peripheral neuropathy, central nervous system vasculitis), and reduced complement levels ([Agnello, 1992](#)). Glomerular disease results from deposition of HCV-related immune complexes in the glomeruli ([Johnson, 1993](#)). Because patients with chronic hepatitis C frequently have laboratory evidence of cryoglobulins (>50% in some series), antiviral treatment is imperative for those with the syndrome of cryoglobulinemia and symptoms or objective

evidence of end-organ manifestations. Limited data with DAA therapy in the setting of vasculitis end-organ disease related to cryoglobulinemia have demonstrated responses in 20% to 90% of patients ([Comarmond, 2017](#)); ([Emery, 2017](#)). Despite this, patients with severe end-organ disease may still require treatment with plasmapheresis or rituximab ([Emery, 2017](#)).

Diabetes

The relationship between chronic hepatitis C and diabetes (most notably type 2 diabetes and insulin resistance) is complex and incompletely understood. The prevalence and incidence of diabetes is increased in the context of hepatitis C ([White, 2008](#)). In the United States, type 2 diabetes occurs more frequently in HCV-infected patients, with a >3-fold greater risk in persons older than 40 years ([Mehta, 2000](#)). The positive correlation between plasma HCV RNA load and established markers of insulin resistance confirms this relationship ([Yoneda, 2007](#)). Insulin resistance and type 2 diabetes are independent predictors of accelerated liver fibrosis progression ([Petta, 2008](#)). Patients with type 2 diabetes and insulin resistance are also at increased risk for HCC ([Hung, 2010](#)).

Successful antiviral treatment has been associated with improved markers of insulin resistance and a greatly reduced incidence of new-onset type 2 diabetes and insulin resistance in HCV-infected patients ([Arase, 2009](#)). Most recently, HCV antiviral therapy has been shown to improve clinical outcomes related to diabetes. In a large prospective cohort from Taiwan, the incidence rates of end-stage renal disease, ischemic stroke, and acute coronary syndrome were greatly reduced in HCV-infected patients with diabetes who received antiviral therapy compared to untreated, matched controls ([Hsu, 2014](#)). Therefore, antiviral therapy may prevent progression to diabetes in HCV-infected patients with prediabetes, and may reduce renal and cardiovascular complications in HCV-infected patients with established diabetes.

Fatigue

Fatigue is the most frequently reported symptom in patients with chronic hepatitis C, and has a major effect on quality of life and activity level as evidenced by numerous measures of impaired quality of life ([Foster, 1998](#)). The presence and severity of fatigue appears to correlate poorly with disease activity, although it may be more common and severe in HCV-infected individuals with cirrhosis ([Poynard, 2002a](#)). Despite difficulties in separating fatigue symptoms associated with hepatitis C from those associated with other concurrent conditions (eg, anemia, depression), numerous studies have reported a reduction in fatigue after cure of HCV infection ([Bonkovsky, 2007](#)). In the Virahep-C study, 401 patients with HCV infection were evaluated for fatigue prior to and after treatment, using validated scales to assess the presence and severity of fatigue ([Sarkar, 2012](#)). At baseline, 52% of patients reported having fatigue, which was more frequent and severe in patients with cirrhosis than in those without cirrhosis. Achieving SVR was associated with a substantial decrease in the frequency and severity of fatigue.

A recent analysis of 413 patients from the NEUTRINO and FUSION trials who were treated with a sofosbuvir-containing regimen and achieved SVR12 demonstrated improvement in patient fatigue (present in 12%) from the pretreatment level ([Younossi, 2014](#)). After achieving SVR12, participants had marked improvements in fatigue over their pretreatment scores, measured by 3 separate validated questionnaires. Additional studies support and extend these findings beyond fatigue, with improvements in overall health-related quality of life and work productivity observed following successful HCV therapy ([Gerber, 2016](#)); ([Younossi, 2015b](#)); ([Younossi, 2015c](#)); ([Younossi, 2015d](#)); ([Younossi, 2015e](#)); ([Younossi, 2016a](#)).

Dermatologic Manifestations

The reported prevalence of HCV infection in patients with porphyria cutanea tarda approximates 50% and occurs disproportionately in those with cirrhosis ([Gisbert, 2003](#)). The treatment of choice for active porphyria cutanea tarda is iron reduction by phlebotomy and maintenance of a mildly iron-reduced state without anemia. Although improvement of porphyria cutanea tarda during HCV treatment with interferon has frequently been described ([Takikawa, 1995](#)), there are currently insufficient data to determine whether HCV DAA therapy and achievement of SVR results in porphyria cutanea tarda improvement.

Lichen planus is characterized by pruritic papules involving mucous membranes, hair, and nails. HCV antibodies are present in 10% to 40% of patients with lichen planus but a causal link with chronic HCV infection is not established. Resolution of lichen planus has been reported with interferon-based regimens, but there have also been reports of exacerbation with these treatments. Although it is unknown whether DAAs will have more success against lichen planus,

treatment with interferon-free regimens would appear to be a more advisable approach to addressing this disorder ([Gumber, 1995](#)); ([Sayiner, 2017](#)).

Benefit of Treatment to Reduce Transmission

Persons who have successfully achieved SVR (virologic cure) no longer transmit the virus to others. As such, successful treatment of HCV infection benefits public health. Several health models have shown that even modest increases in successful treatment of HCV infection among persons who inject drugs can decrease prevalence and incidence ([Martin, 2013a](#)); ([Durier, 2012](#)); ([Martin, 2013b](#)); ([Hellard, 2012](#)); ([Harris, 2016](#)). Models developed to estimate the impact of HCV testing and treatment on the burden of hepatitis C at a country level reveal that large decreases in HCV prevalence and incidence are possible as more persons are successfully treated ([Wedemeyer, 2014](#)).

There are also benefits to eradicating HCV infection between couples and among families, thus eliminating the perception that an individual might be contagious. In addition, mother-to-child transmission of HCV does not occur if the woman is not viremic, providing an additional benefit of curing a woman before she becomes pregnant ([Thomas, 1998](#)). The safety and efficacy of treating women who are already pregnant, however, to prevent transmission to the fetus have not yet been established. Thus, treatment is not recommended for pregnant women.

The Society for Healthcare Epidemiology of America (SHEA) advises that healthcare workers who have substantial HCV viral replication ($\geq 10^4$ genome equivalents/mL) be restricted from performing procedures that are prone to exposure ([Henderson, 2010](#)) and that all healthcare workers with confirmed chronic HCV infection should be treated. For reasons already stated, the achievement of SVR in such individuals will not only eliminate the risk of HCV transmission to patients but also decrease circumstantial loss of experienced clinicians. Given concerns about underreporting of infection and transmission ([Henderson, 2010](#)), the availability of effective, all-oral regimens should lead to greater willingness on the part of exposure-prone clinicians to be tested and treated.

Successful treatment of HCV-infected persons at greatest risk for transmission represents a formidable tool to help stop HCV transmission in those who continue to engage in high-risk behaviors. To guide implementation of hepatitis C treatment as a prevention strategy, studies are needed to define the best candidates for treatment to stop transmission, the additional interventions needed to maximize the benefits of HCV treatment (eg, preventing reinfection), and the cost-effectiveness of the strategies when used in target populations.

Persons Who Inject Drugs

Injection drug use (IDU) is the most common risk factor for HCV infection in the United States and Europe, with an HCV seroprevalence rate of 10% to 70% ([Amon, 2008](#)); ([Nelson, 2011](#)). IDU also accounts for the majority of new HCV infections (approximately 70%) and is the key driving force in the perpetuation of the epidemic. Given these facts and the absence of an effective vaccine against HCV, testing and linkage to care combined with treatment of HCV infection with potent DAAs has the potential to dramatically decrease HCV incidence and prevalence ([Martin, 2013b](#)). However, treatment-based strategies to prevent HCV transmission have yet to be studied, including how to integrate hepatitis C treatment with other risk-reduction strategies (eg, opiate substitution therapy, and needle and syringe exchange programs) ([Martin, 2013a](#)).

In studies of interferon-based treatments in persons who inject drugs, adherence and efficacy rates are comparable to those of patients who do not use injected drugs. A meta-analysis of treatment with peginterferon, with or without ribavirin, in active or recent injection drug users showed SVR rates of 37% and 67% for genotype 1 or 4, and 2 or 3, respectively ([Aspinall, 2013](#)). With the introduction of shorter, better-tolerated, and more efficacious interferon-free therapies, these SVR rates are expected to improve. Importantly, the rate of reinfection in this population is lower (2.4/100 person-years of observation) than that of incident infection in the general population of injection drug users (6.1 to 27.2/100 person-years), although reinfection increases with active or ongoing IDU (6.44/100 person-years) and available data on follow-up duration are limited ([Aspinall, 2013](#)); ([Grady, 2013](#)).

Ideally, treatment of HCV-infected persons who inject drugs should be delivered in a multidisciplinary care setting with services to reduce the risk of reinfection and for management of the common social and psychiatric comorbidities in this

population ([Murphy 2015](#)); ([Dore, 2016](#)); ([Mathei 2016](#)); ([Midgard 2016](#)). Regardless of the treatment setting, recent or active IDU should not be seen as an absolute contraindication to HCV therapy. There is strong evidence from various settings in which persons who inject drugs have demonstrated adherence to treatment and low rates of reinfection, countering arguments that have been commonly used to limit treatment access in this patient population ([Aspinall, 2013](#)); ([Hellard, 2014](#)); ([Grebely, 2011](#)). Indeed, combining HCV treatment with needle exchange and opioid agonist therapy programs in this population with a high prevalence of HCV has shown great value in decreasing the burden of HCV disease. Elegant modeling studies illustrate high return on the modest investment of addressing this often-ignored segment of the HCV-infected population ([Martin, 2013b](#)). These conclusions were drawn before the introduction of the latest DAA regimens. Conversely, there are no data to support the utility of pretreatment screening for illicit drug or alcohol use in identifying a population more likely to successfully complete HCV therapy. These requirements should be abandoned because they create barriers to treatment, add unnecessary cost and effort, and potentially exclude populations that are likely to obtain substantial benefit from therapy. Scaling up HCV treatment in persons who inject drugs is necessary to positively impact the HCV epidemic in the US and globally.

HIV-Infected Men Who Have Sex With Men

Since 2000, a dramatic increase in incident HCV infections among HIV-infected men who have sex with men (MSM) who did not report IDU as a risk factor has been demonstrated in several US cities ([van de Laar, 2010](#)); ([Samandari, 2017](#)). Recognition and treatment of HCV infection (including acute infection) in this population may represent an important step in preventing subsequent infections ([Martin, 2016](#)). As with persons who inject drugs, HIV/HCV-coinfected MSM who engage in ongoing high-risk sexual practices should be treated for their HCV infection in conjunction with continued education about risk-reduction strategies. In particular, safer-sex strategies should be emphasized given the high rate of reinfection after SVR, which may approach 30% over 2 years in HIV-infected MSM with acute HCV infection ([Lambers, 2011](#)).

Some of the best examples of HCV treatment as prevention of transmission have come from well characterized cohorts of HIV/HCV coinfecting MSM. In the Dutch acute HCV in HIV study (DAHHS) cohort, a 51% decrease in HCV incidence among MSM living with HIV was realized in just 2 years after implementing a comprehensive HCV screening and immediate treatment program ([Boerekamps, 2017](#)). Similarly, in the Swiss HIV cohort study (SHCS), a 92.5% reduction in HCV prevalence and 51% decrease in incident HCV infections was realized shortly after implementing universal screening and treatment within an MSM cohort living with HIV ([Braun, 2018](#)).

Incarcerated Persons

Among incarcerated individuals, the rate of HCV seroprevalence ranges from 30% to 60% ([Post, 2013](#)) and the rate of acute infection is approximately 1% ([Larney, 2013](#)). Screening for HCV infection is relatively uncommon in state prison systems. Treatment uptake has historically been limited, in part because of the toxic effects and long treatment duration of older interferon-based therapies as well as cost concerns ([Spaulding, 2006](#)). In particular, truncation of HCV treatment owing to release from prison has been cited as a major limitation to widespread, effective HCV treatment in correctional facilities ([Post, 2013](#)); ([Chew, 2009](#)). Shorter HCV treatment duration with DAA regimens reduces stay-related barriers to HCV treatment in prisons. Likewise, the improved safety of DAA regimens diminishes concerns about toxic effects. Coordinated treatment efforts within prison systems would likely rapidly decrease HCV prevalence in this at-risk population ([He, 2016](#)), although research is needed in this area.

Persons on Hemodialysis


HCV prevalence is markedly elevated in persons on hemodialysis, ranging from 2.6% to 22.9% in a large multinational study ([Fissell, 2004](#)). US studies found a similarly elevated prevalence of 7.8% to 8.9% ([CDC, 2001](#)); ([Finelli, 2005](#)). Importantly, the seroprevalence of HCV was found to increase with time on dialysis, suggesting that nosocomial transmission, among other risk factors, plays a role in HCV acquisition in these patients ([Fissell, 2004](#)). Improved education and strict adherence to universal precautions can drastically reduce nosocomial HCV transmission risk for persons on hemodialysis ([Jadoul, 1998](#)), but clearance of HCV viremia through treatment-induced SVR eliminates the potential for transmission.

HCV-infected persons on hemodialysis have a decreased quality of life and increased mortality compared to those who are uninfected ([Fabrizi, 2002](#)); ([Fabrizi, 2007](#)); ([Fabrizi, 2009](#)). HCV infection in this population also has a deleterious impact on kidney transplantation outcomes with decreased patient and graft survival ([Fabrizi, 2014](#)). The increased risk for nosocomial transmission and the substantial clinical impact of HCV infection in those on hemodialysis are compelling arguments for HCV therapy as effective antiviral regimens that can be used in persons with advanced renal failure are now available (see [Patients with Renal Impairment](#)).

Patients Unlikely to Benefit From HCV Treatment

Patients with a limited life expectancy that cannot be remediated by HCV treatment, liver transplantation, or another directed therapy do not require antiviral treatment. Patients with a short life expectancy owing to liver disease should be managed in consultation with an expert. Chronic hepatitis C is associated with a wide range of comorbid conditions ([Butt, 2011](#)); ([Louie, 2012](#)). Little evidence exists to support initiation of HCV treatment in patients with a limited life expectancy (<12 months) owing to nonliver-related comorbid conditions. For these patients, the benefits of HCV treatment are unlikely to be realized and palliative care strategies should take precedence ([Holmes, 2006](#)); ([Maddison, 2011](#)).

Pretreatment Assessment

Recommendation for Pretreatment Assessment	
RECOMMENDED	RATING 
Evaluation for advanced fibrosis using noninvasive markers and/or elastography, and rarely liver biopsy, is recommended for all persons with HCV infection to facilitate decision making regarding HCV treatment strategy and determine the need for initiating additional measures for the management of cirrhosis (eg, hepatocellular carcinoma screening) (see HCV Testing and Linkage to Care).	I, A

An accurate assessment of fibrosis remains vital as the degree of hepatic fibrosis is one of the most robust prognostic factors used to predict HCV disease progression and clinical outcomes ([Everhart, 2010](#)). Individuals with severe fibrosis require surveillance monitoring for liver cancer, esophageal varices, and hepatic function ([Garcia-Tsao, 2007](#)); ([Bruix, 2011](#)). In some instances, the recommended duration of treatment is also longer.

Although liver biopsy is the diagnostic standard, sampling error and observer variability limit test performance, particularly when inadequate sampling occurs. Up to 1/3 of bilobar biopsies had a difference of at least 1 stage between the lobes ([Bedossa, 2003](#)). In addition, the test is invasive and minor complications are common, limiting patient and practitioner acceptance. Although rare, serious complications such as bleeding are well recognized.


Noninvasive tests to stage the degree of fibrosis in patients with chronic HCV infection include models incorporating indirect serum biomarkers (routine tests), direct serum biomarkers (components of the extracellular matrix produced by activated hepatic stellate cells), and vibration-controlled transient liver elastography. No single method is recognized to have high accuracy alone, and each test must be interpreted carefully. A publication from the Agency for Healthcare Research and Quality found evidence in support of a number of blood tests; however, at best, they are only moderately useful for identifying clinically significant fibrosis or cirrhosis ([Selph, 2014](#)).

Vibration-controlled transient liver elastography is a noninvasive way to measure liver stiffness and correlates well with measurement of substantial fibrosis or cirrhosis in patients with chronic HCV infection. The measurement range, however, overlaps between stages ([Ziol, 2005](#)); ([Afdhal, 2015](#)); ([Castera, 2005](#)).

The most efficient approach to fibrosis assessment is to combine direct biomarkers and vibration-controlled transient liver elastography ([Boursier, 2012](#)); ([European Association for the Study of the Liver and Asociacion Latinoamericana para el Estudio del Hgado, 2015](#)). A biopsy should be considered for any patient who has discordant results between the 2 modalities that would affect clinical decision making (eg, one shows cirrhosis and the other does not). The need for liver biopsy with this approach is markedly reduced.

Alternatively, if direct biomarkers or vibration-controlled transient liver elastography are not available, the AST-to-platelet ratio index (APRI) or FIB-4 index score can prove helpful—although neither is sensitive enough to rule out substantial fibrosis ([Sebastiani, 2009](#)); ([Castera, 2010](#)); ([Chou, 2013](#)). Biopsy should be considered for those in whom more accurate fibrosis staging would impact treatment decisions. Individuals with clinically evident cirrhosis do not require additional staging (biopsy or noninvasive assessment).

Recommendation for Repeat Liver Disease Assessment

RECOMMENDED	RATING 
Ongoing assessment of liver disease is recommended for persons in whom therapy is deferred.	I, C

Ongoing assessment of liver disease is especially important in patients for whom therapy has been deferred. In line with evidence-driven recommendations for treatment of nearly all HCV-infected patients, several factors must be taken into consideration if treatment deferral is entertained or mandated by lack of medication access. As noted, strong and accumulating evidence argue against deferral because of decreased all-cause morbidity and mortality, prevention of onward transmission, and quality-of-life improvements for patients treated regardless of baseline fibrosis. Additionally, successful HCV treatment may improve or prevent extrahepatic complications, including diabetes mellitus, cardiovascular disease, renal disease, and B-cell non-Hodgkin lymphoma ([Conjeevaram, 2011](#)); ([Hsu, 2015](#)); ([Torres, 2015](#)), which are not tied to fibrosis stage ([Allison, 2015](#)); ([Petta, 2016](#)). Deferral practices based on fibrosis stage alone are inadequate and shortsighted.

Fibrosis progression varies markedly between individuals based on host, environmental, and viral factors (Table 1); ([Feld, 2006](#)). Fibrosis may not progress linearly. Some individuals (often those aged >50 years) may progress slowly for many years followed by accelerated fibrosis progression. Others may never develop substantial liver fibrosis despite longstanding infection. The presence of existing fibrosis is a strong risk factor for future fibrosis progression. Fibrosis results from chronic hepatic necroinflammation; thus, a higher activity grade on liver biopsy and higher serum transaminase levels are associated with more rapid fibrosis progression ([Ghany, 2003](#)). However, even patients with a normal ALT level may develop substantial liver fibrosis over time ([Pradat, 2002](#)); ([Nutt, 2000](#)). The limitations of transient elastography and liver biopsy in ascertaining the progression of fibrosis must be recognized.

Host factors associated with more rapid fibrosis progression include male sex, longer duration of infection, and older age at the time of infection ([Poynard, 2001](#)). Many patients have concomitant nonalcoholic fatty liver disease. The presence of hepatic steatosis (with or without steatohepatitis) on liver biopsy, elevated body mass index, insulin resistance, and iron overload are associated with fibrosis progression ([Konerman, 2014](#)); ([Everhart, 2009](#)). Chronic alcohol use is an important risk factor because alcohol consumption has been associated with more rapid fibrosis progression ([Feld, 2006](#)). A safe amount of alcohol consumption has not been established. Cigarette smoking may also lead to more rapid fibrosis progression. For more counseling recommendations, see [Testing and Linkage to Care](#).

Immunosuppression leads to more rapid fibrosis progression, particularly in the settings of HIV/HCV coinfection and solid organ transplantation ([Macias, 2009](#)); ([Konerman, 2014](#)); ([Berenguer, 2013](#)). Therefore, immunocompromised patients should be treated even if they have mild liver fibrosis at presentation.

HCV RNA level does not correlate with stage of disease (degree of inflammation or fibrosis). Available data suggest that fibrosis progression occurs most rapidly in patients with genotype 3 ([Kanwal, 2014](#)); ([Bochud, 2009](#)). Aside from coinfection with HBV or HIV, no other viral factors are consistently associated with disease progression.

Although an ideal interval for assessment has not been established, annual evaluation is appropriate to discuss modifiable risk factors and update testing for hepatic function and markers of disease progression. For all individuals with advanced fibrosis, liver cancer screening dictates a minimum of evaluation every 6 months.

Table. Factors Associated With Accelerated Fibrosis Progression

Host	Viral
<p>Nonmodifiable</p> <ul style="list-style-type: none"> • Fibrosis stage • Inflammation grade • Older age at time of infection • Male sex • Organ transplant <p>Modifiable</p> <ul style="list-style-type: none"> • Alcohol consumption • Nonalcoholic fatty liver disease • Obesity • Insulin resistance 	<ul style="list-style-type: none"> • Genotype 3 • Coinfection with hepatitis B virus or HIV

Last update: November 6, 2019

Overview of Cost, Reimbursement, and Cost-Effectiveness Considerations for Hepatitis C Treatment Regimens

The hepatitis C guidance describes diagnosis, linkage to care, and treatment for people with HCV infection ([AASLD/IDSA, 2020](#)). Reduced access to treatment, however, is a common challenge due to restrictions on drug reimbursement. This section summarizes the US payer system, explains the concepts of cost, price, cost-effectiveness, value, and affordability, and addresses the cost-effectiveness of HCV treatment. Although these terms may sound similar, the following discussion seeks to clarify them regarding HCV therapy. This section aims to be informational. As explained, actual costs are rarely known. Accordingly, the HCV guidance does not currently utilize cost-effectiveness analysis to guide recommendations.

Drug Cost and Reimbursement

Many organizations are involved with hepatitis C drug distribution and each can impact costs as well as decisions about which regimens are reimbursed ([US GAO, 2015](#)); ([US CBO, 2015](#)). The roles these organizations have in determining the actual price paid for drugs and who has access to treatment include the following:

- Pharmaceutical companies determine the wholesale acquisition cost (WAC) of a drug (analogous to a sticker price). The company negotiates contracts with other organizations within the pharmaceutical supply chain that allow for rebates or discounts to decrease the actual price paid.
- Pharmacy benefit managers (PBMs) act as intermediaries between pharmaceutical companies and health

insurance companies. They negotiate contracts that may include restrictions on the types of providers or patients who can be reimbursed for treatment. They might also offer exclusivity (restrictions on which medications can be prescribed) in exchange for lower negotiated prices, often provided in the form of WAC discounts.

- Private insurance companies often have separate pharmacy and medical budgets, and use PBMs or directly negotiate drug pricing with pharmaceutical companies. Insurance companies determine formulary placement, which impacts the choice of regimens and out-of-pocket expenses for patients. An insurance company can cover private, managed care Medicaid, and Medicare plans and have different formularies for each line of business.
- Medicaid is a heterogeneous consortium of insurance plans that includes fee-for-service and managed care options. Most plans negotiate rebates with pharmaceutical manufacturers (through PBMs or individually). For single-source drugs such as all-oral HCV treatments, Medicaid plans receive the lowest price offered to any other payer (outside of certain government agencies), and the minimum Medicaid drug rebate is 23.1% of the average manufacturer price (AMP). Differences in negotiated contracts between plans have led to Medicaid patients in different states having widely varied access to HCV therapy ([Barua, 2015](#)); ([Canary, 2015](#)); ([Lo Re, 2016](#)). State Medicaid programs have benefited from the Patient Protection and Affordable Care Act (ACA), although such benefits are mitigated in states that have opted out of expanding Medicaid coverage under the ACA. As the price of HCV therapies has decreased, states have loosened their Medicaid treatment restrictions with a growing number providing treatment to all infected persons. Many states, however, continue to restrict access to HCV treatment based on stage of liver fibrosis or history of recent drug use. Proposed rollbacks of Medicaid expansion implemented under the ACA threaten to reduce insurance coverage among HCV-infected people and could lead to new treatment restrictions.
- Medicare covers HCV drugs through part D benefits and is prohibited by law from directly negotiating drug prices. These drug plans are offered through PBMs or commercial health plans, which may negotiate discounts or rebates with pharmaceutical companies.
- The Veterans Health Administration receives mandated rebates through the Federal Supply Schedule program, which sets drug prices for several government agencies (including the Department of Veterans Affairs, federal prisons, and the Department of Defense) and typically receives substantial discounts over average wholesale price (AWP).
- State prisons and jails are usually excluded from Medicaid-related rebates and often do not have the negotiating leverage of larger organizations and, therefore, may pay higher prices than most other organizations.
- Specialty pharmacies receive dispensing fees and may receive additional payments from contracted insurance companies, PBMs, or pharmaceutical companies to provide services such as adherence support and/or management of adverse effects, and outcome measurements, such as early discontinuation rates and sustained virologic response rates.
- Patients incur costs (eg, copayment or coinsurance) determined by their pharmacy plan. Patient assistance programs offered by pharmaceutical companies or foundations can cover many of these out-of-pocket expenses or provide drugs at no cost to qualified patients who are unable to pay.

Except for mandated rebates, negotiated drug prices are considered confidential business contracts. Therefore, there is almost no transparency regarding the actual prices paid for hepatitis C drugs ([Saag, 2015](#)). However, the average negotiated discount of 22% in 2014 increased to 46% less than the WAC in 2015, implying that many payers are paying well below the WAC for HCV medications ([Committee on Finance US Senate, 2016](#)).

Cost-Effectiveness

Cost-effectiveness analysis (CEA) compares the relative costs and outcomes of 2 or more interventions. CEA explicitly recognizes budget limitations for healthcare spending and seeks to maximize public health benefits within those budgetary constraints. The core question that CEA addresses is whether to invest limited healthcare dollars in a new treatment/therapy or use that money to invest in another healthcare intervention that would provide better outcomes for the same monetary investment. The focus of CEA is, therefore, not simply cost or saving money but health benefits. It assumes that all available resources will be spent and provides a framework for prioritizing among available treatment options by formally assessing the comparative costs and health benefits accrued from a new treatment relative to current treatment.

The cost-effectiveness of a treatment is typically expressed as an incremental cost-effectiveness ratio (ICER).

$$\frac{\text{cost new treatment} - \text{cost current treatment}}{\text{benefit new treatment} - \text{benefit current treatment}}$$

Estimating and Interpreting the ICER

Estimating and interpreting the ICER requires answering 3 questions:

1. *How much more money will be spent with the new treatment versus the old treatment?*

The additional cost of new treatment includes that of new medications as well as the costs that will be avoided by preventing disease complications. Prevention of long-term complications is especially important when considering the cost-effectiveness of HCV treatments because the costs of the therapy are immediate, while those avoided by preventing advanced liver disease and other complications of chronic infection often accrue years in the future.

2. *How much more benefit will occur with the new versus the old treatment?*

Life expectancy is a valuable measure of benefit but considering only mortality benefits fails to recognize the value of treatments that improve quality of life. The quality-adjusted life-year (QALY) provides a measure that integrates both longevity and quality of life and is the preferred outcome for CEA.

3. *How is the ICER to be interpreted?*

The ideal CEA would list every possible healthcare intervention, its lifetime medical cost, and QALYs lived. Such a list would allow for perfect theoretical prioritization of spending to maximize QALY across the population. In reality, CEA compares the ICER for a specific treatment to a threshold value and rejects treatments with an ICER exceeding a particular threshold as not being cost-effective. The threshold value is referred to as the societal willingness-to-pay threshold. It is not meant to be a valuation of how much society is willing to pay to save a life. Rather, it is meant to reflect the average return in QALY expected if the available budget was not used to provide a new treatment but instead invested into the current healthcare system. In the United States, the willingness-to-pay threshold is typically considered to be \$50,000 or \$100,000 per QALY gained.

Affordability

An intervention that is cost-effective is not necessarily affordable. Affordability refers to whether a payer has sufficient resources in its annual budget to pay for a new therapy for all who might need or want it within that year. Several characteristics of CEA limit its ability to speak to the budgetary impact of interventions being implemented in the real world.

1. *Perspective on cost*

CEA seeks to inform decisions about how society should prioritize healthcare spending. As such, it typically assumes a societal perspective on costs and includes all costs from all payers, including out-of-pocket expenses for the patient. When making coverage decisions for therapy, however, an insurer considers only its own revenues and expenses.

2. *Time horizon*

From a societal perspective, CEA uses a lifetime time horizon, meaning it considers lifetime costs and benefits, including those that occur in the distant future. Business budget planning, however, typically assumes a 1-year to 5-year perspective. Savings that may accrue 30 years from now have no impact on spending decisions today because they have little bearing on the solvency of the current budget.

3. *Weak association between willingness-to-pay and the real-world bottom line*

Societal willingness-to-pay thresholds in CEAs are not based on actual budget calculations and have little

relationship to a payer's bottom line. Willingness-to-pay is meant to be an estimate of the opportunity cost of investing in a new therapy. In economics, opportunity cost refers to how else that money could have been spent and the benefits lost from not investing in that alternative ([Wong, 2017a](#)). When payers make a decision about coverage, the calculation is more straightforward and relates to the short-term cost of medications and the budgetary impact. Given the rapid development of new technologies and therapies, funding all of them (even if they all fell below the societal willingness-to-pay threshold) would likely lead to uncontrolled growth in demand and exceed the limited healthcare budget.

There is no formula that provides a good means of integrating the concerns of value and affordability. When new HCV therapies are deemed cost-effective, it indicates that these therapies provide good benefit for the resources invested and providing such therapy to more people would be a good long-term investment. Determining the total resources that can be spent on HCV treatment, however, depends on political and economic factors that are not captured by cost-effectiveness determinations.

Cost-Effectiveness of Current Direct-Acting Antiviral Regimens for Hepatitis C Treatment

Since the first direct-acting antivirals (DAAs) received US Food and Drug Administration approval in 2011, several cost-effectiveness investigations have compared DAA-based regimens to previous standard-of-care regimens to calculate ICERs. They have also investigated the cost-effectiveness of eliminating HCV treatment restrictions. Compared to interferon-based regimens, the ICER for DAAs has consistently been estimated at <\$100,000 per QALY for all genotypes and fibrosis stages.

Several studies have compared DAA regimens against one another. In general, when given a choice between recommended HCV DAA regimens, the less costly regimen is preferred as a more efficient use of resources (even if it requires multiple tablet dosing). Because of the similar efficacy of most DAA regimens, cost becomes the critical factor driving relative cost-effectiveness. Studies have also estimated the cost-effectiveness of HCV treatment in special populations, including patients awaiting liver transplantation, HIV/HCV-coinfected patients, those with chronic kidney disease, persons who inject drugs, and adolescents—all with favorable ICERs. At this time, it is reasonable to conclude that DAA regimens provide good value for the resources invested.

Cost vs Affordability for HCV Treatment

Despite a growing body of evidence that HCV treatment is cost-effective and may even be cost saving over the long term in some cases, many US payers—especially those offering Medicaid insurance products—continue to limit access to HCV treatment. Access has improved as cost has decreased but limitations remain. Proposed reductions in healthcare spending for Medicaid would likely exacerbate the problem as the value of the HCV medications would remain unchanged but the resources available to provide them would shrink.

Cost-Effectiveness of Screening for HCV

Several cost-effectiveness studies demonstrate that routine, one-time testing for HCV among all adults in the US would likely identify a substantial number of cases of HCV that are currently being missed, and that doing so would be cost-effective. One study employed simulation modeling to compare several versions of routine guidance, including routine testing for adults over the ages of 40 years, 30 years, and 18 years and found that routine testing for all adults aged ≥ 18 years was cost-effective compared to risk-based screening, and potentially cost-saving compared to testing only those aged ≥ 30 years or aged ≥ 40 years ([Barocas, 2018](#)). The study further found that routine testing remained cost-effective unless HCV infection had no impact on healthcare utilization and no impact on quality of life. Another research group similarly found that routine testing of all adults aged ≥ 18 years is likely cost-effective compared to risk-based screening, so long as the prevalence of HCV among those born after 1965 exceeds 0.07% ([Eckman, 2019](#)). Notably, these studies reached similar conclusions despite being conducted entirely independently and employing different simulation modeling approaches. Further, a variety of studies have examined the cost-effectiveness of routine HCV testing in specific venues, including correctional settings ([He, 2016](#)), prenatal care settings ([Chaillon, 2019](#)); ([Tasillo, 2019](#)), substance use treatment centers ([Schackman, 2018](#)); ([Schackman, 2015](#)), and federally qualified health centers ([Assoumou, 2018](#)). All of them found that routine testing and treatment for HCV was cost-effective, even when linkage to HCV treatment after testing was poor, and even when the rate of HCV reinfection among injection drug users is common.

Generally, routine HCV testing is cost-effective because the incidence and prevalence of HCV remain high in people who inject drugs with a notable rising prevalence in young adults who may not readily report their stigmatized risk behaviors. Studies conducted in urban emergency departments in the US, for example, reveal that 15% to 25% of patients with previously unidentified HCV infection were born after 1965 and/or have no reported history of injection drug use and are, therefore, missed by even perfect implementation of risk-based screening ([Schechter-Perkins, 2018](#)); ([Hsieh, 2016](#)); ([Lyons, 2016](#)). Reinfection among those actively using drugs is common but because screening is a low-cost intervention, and therapy is both highly effective and cost-effective, routine testing provides good economic value (ie, cost-effective) even when many people need to be tested and treated more than once during their lifetime.

Conclusions


Many studies have demonstrated the economic value of HCV screening ([Chaillon, 2019](#)); ([Eckman, 2019](#)); ([Tasillo, 2019](#)); ([Assoumou, 2018](#)); ([Barocas, 2018](#)); ([Schackman, 2018](#)); ([Schechter-Perkins, 2018](#)); ([Lyons, 2016](#)); ([Hsieh, 2016](#)); ([Schackman, 2015](#)) and treatment ([Goel, 2018](#)); ([Chhatwal, 2017](#)); ([He, 2017](#)); ([Chahal, 2016](#)); ([Chhatwal, 2015](#)); ([Chidi, 2016](#)); ([Martin, 2016a](#)); ([Lin, 2015](#)); ([Najafzadeh, 2015](#)); ([Rein, 2015](#)); ([Tice, 2015](#)); ([Younossi, 2015a](#)) and made it clear that HCV screening and therapy are cost-effective. In response, in 2020, both the US Centers for Disease Control (CDC) and Prevention and the US Preventive Service Task Force (USPSTF) recommended routine, one-time HCV testing for all US adults aged ≥ 18 years without effort to identify risk factors, as well as repeat testing for those with ongoing risk factors ([Owens, 2020](#)); ([Schillie, 2020](#)). The high cost of HCV medications and the high prevalence of disease have led to limited access for some patients. The issue is complex. Although the wholesale acquisition costs of HCV drugs often make treatment appear unaffordable, the reality is that insurers, PBMs, and government agencies negotiate pricing and few actually pay this much-publicized price. Negotiated pricing and cost structure for pharmaceutical products in the US are not transparent, however. Thus, it is difficult to estimate the true budgetary impact of providing HCV drugs. Competition and negotiated pricing have reduced prices substantially but cost continues to limit the public health impact of DAA therapies. Insurers, government, and pharmaceutical companies should work together to bring medication prices to the point where all persons in need of treatment are able to afford and readily access it.

Last update: August 27, 2020

Monitoring Patients Who Are Starting HCV Treatment, Are on Treatment, or Have Completed Therapy

This section provides guidance on monitoring patients with chronic hepatitis C virus (HCV) infection who are starting direct-acting antiviral (DAA) treatment, are on treatment, or have completed therapy. It is divided into 4 parts: pretreatment and on-treatment monitoring; post-treatment follow-up for persons in whom treatment failed to clear the virus; post-treatment follow-up for those who achieve a sustained virologic response (SVR; virologic cure); and additional considerations if treatment includes ribavirin.

Pretreatment and On-Treatment Monitoring

Recommended Assessments Prior to Starting DAA Therapy	
RECOMMENDED	RATING 
Staging of hepatic fibrosis is essential prior to HCV treatment (see Testing and Linkage to Care and see When and in Whom to Treat).	I, C

Recommended Assessments Prior to Starting DAA Therapy

Assessment of potential drug-drug interactions with concomitant medications is recommended prior to starting DAA therapy and, when possible, an interacting co-medication should be stopped or switched to an alternative with less risk for potential interaction during HCV treatment. (See [Table of Drug Interactions with Direct-Acting Antivirals and Selected Concomitant Medications](#) below or use an online resource such as [University of Liverpool interaction checker](#).)

Patients should be educated about the proper administration of medications (eg, dose, frequency of medicines, food effect, missed doses, adverse effects, etc), the crucial importance of adherence, and the need to inform the healthcare provider about any changes to their medication regimen.

The following laboratory tests are recommended within 6 months prior to starting DAA therapy:

- Complete blood count (CBC)
- International normalized ratio (INR)
- Hepatic function panel (ie, albumin, total and direct bilirubin, alanine aminotransferase [ALT], aspartate aminotransferase [AST], and alkaline phosphatase levels)
- Calculated glomerular filtration rate (eGFR)

The following laboratory tests are recommended anytime prior to starting DAA therapy:

- Quantitative HCV RNA (HCV viral load)
- If a non-pan-genotypic DAA will be prescribed, then test for HCV genotype and subtype.

The safety of ribavirin-free DAA regimens in humans has not been established during pregnancy and for nursing mothers, so counseling should be offered to women of childbearing age before beginning HCV treatment. (See ribavirin pregnancy recommendations below.)

I, C

All patients initiating DAA therapy should be assessed for active hepatitis B virus (HBV) coinfection with HBV surface antigen (HBsAg) testing, and for evidence of prior infection with HBV core antibody (anti-HBc) and HBV surface antibody (anti-HBs) testing.

IIa, B

Patients found or known to be HBsAg-positive should be assessed for whether their HBV DNA level meets [AASLD criteria for HBV treatment and initiation of antiviral therapy for HBV](#).

Strong, Moderate^a

All patients initiating DAA therapy should be assessed for HIV coinfection.

IIa, B

Testing for the presence of resistance-associated substitutions (RASs) prior to starting treatment should be performed as recommended in the [Initial Treatment](#) and the [Retreatment](#) sections. Additional information about RAS testing can be found in the [HCV Resistance Primer](#).

IIb, B

Patients scheduled to receive an HCV NS3 protease inhibitor (ie, grazoprevir, voxilaprevir, glecaprevir) should be assessed for a history of decompensated liver disease and liver disease severity using the Child-Turcotte-Pugh (CTP) score (see third-party calculator).


I, A

- Patients with current or prior history of decompensated liver disease or with a current CTP score ≥ 7 should **not** receive treatment with regimens that contain NS3 protease inhibitors due to increased blood levels and/or lack of safety data.

Recommended Assessments Prior to Starting DAA Therapy

^a Unlike the AASLD/IDSA HCV guidance, the AASLD guidelines for treatment of chronic hepatitis B uses the GRADE approach to rate recommendations; please see that [document](#) for further information about this rating system.

Recommended Monitoring During Antiviral Therapy

RECOMMENDED	RATING 
Clinic visits or telephone contact are recommended as clinically indicated during treatment to ensure medication adherence, and to monitor for adverse events and potential drug-drug interactions (see table of Drug Interactions with Direct-Acting Antivirals and Selected Concomitant Medications below), especially with newly prescribed medications.	I, B
Inform patients taking diabetes medication of the potential for symptomatic hypoglycemia. On-treatment and post-treatment monitoring for hypoglycemia is recommended.	I, C
Inform patients taking warfarin of the potential for changes in their anticoagulation status. On-treatment and post-treatment INR monitoring for subtherapeutic anticoagulation is recommended.	I, C
Patients receiving elbasvir/grazoprevir should be monitored with a hepatic function panel at 8 weeks and again at 12 weeks if receiving 16 weeks of treatment.	I, B
<p>A ≥ 10-fold increase in ALT values from baseline at any time during treatment should prompt discontinuation of DAA therapy (especially with signs or symptoms of liver inflammation or increasing conjugated bilirubin, alkaline phosphatase, or INR).</p> <p>An increase in ALT < 10-fold from baseline that is accompanied by any weakness, nausea, vomiting, jaundice, or significantly increased bilirubin, alkaline phosphatase, or INR should also prompt discontinuation of DAA therapy.</p> <p>Asymptomatic increases in ALT < 10-fold from baseline should be closely monitored with repeat testing at 2-week intervals. If levels remain persistently elevated, consideration should be given to discontinuation of DAA therapy.</p>	I, B
Quantitative HCV viral load testing is recommended 12 or more weeks after completion of therapy to document SVR (cure).	I, B
<p>For HBsAg-positive patients not already receiving HBV suppressive therapy because their baseline HBV DNA level does not meet treatment criteria, one of two approaches may be taken:</p> <ul style="list-style-type: none"> • Initiate prophylactic HBV antiviral therapy for those with low or undetectable HBV DNA levels. If this course is elected, pending further data, prophylaxis should be continued until 12 weeks after completion of DAA therapy. • Monitor HBV DNA levels monthly during and immediately after DAA therapy. Antiviral treatment for HBV should be given in the event of a rise in HBV DNA > 10-fold above baseline or to > 1000 IU/mL in those with a previously undetectable or unquantifiable HBV DNA level. 	IIa, B

Recommended Monitoring During Antiviral Therapy

The recommended pretreatment testing assumes that a decision to treat with antiviral medications has already been made and that the testing involved in deciding to treat—including testing for HCV genotype and assessment of hepatic fibrosis—has already been completed (see [When and in Whom to Initiate HCV Therapy](#)).

Prior to starting treatment, patients should be evaluated for potential drug-drug interactions with selected antiviral medications by consulting the prescribing information and using other resources (eg, <http://www.hep-druginteractions.org>). The table below lists known drug-drug interactions between HCV DAAs and selected medications.

Table. Drug Interactions with Direct-Acting Antivirals and Selected Concomitant Medications

Concomitant Medications	LDV/SOF	EBR/GZR	VEL/SOF	GLE/PIB	SOF/VEL/VOX
Acid-reducing agents	Antacids H2RA PPI		Antacids H2RA PPI	H2RA PPI	Antacids H2RA PPI
Alpha-1 blockers	Silodosin	Prazosin Silodosin	Prazosin Silodosin	Prazosin Silodosin	Prazosin Silodosin
Antiarrhythmics	Amiodarone Dronedarone	Amiodarone Dronedarone Quinidine	Amiodarone Dronedarone	Amiodarone Dronedarone Quinidine	Amiodarone Dronedarone
	Digoxin Quinidine		Digoxin Quinidine		Digoxin Quinidine
Anticoagulant and antiplatelet agents	Apixaban Dabigatran Edoxaban Rivaroxaban Ticagrelor Warfarin	Apixaban Dabigatran Edoxaban Rivaroxaban Ticagrelor Warfarin	Apixaban Dabigatran Edoxaban Rivaroxaban Ticagrelor Warfarin	Dabigatran	Dabigatran Edoxaban
				Apixaban Edoxaban Rivaroxaban Ticagrelor Warfarin	Apixaban Rivaroxaban Ticagrelor Warfarin
Anticonvulsants and barbiturates	Amobarbital Carbamazepine Eslicarbazine Oxcarbazepine Phenobarbital Phenytoin Primidone Rufinamide	Amobarbital Carbamazepine Eslicarbazine Oxcarbazepine Phenobarbital Phenytoin Primidone	Amobarbital Carbamazepine Eslicarbazine Oxcarbazepine Phenobarbital Phenytoin Primidone Rufinamide	Amobarbital Carbamazepine Eslicarbazine Oxcarbazepine Phenobarbital Phenytoin Primidone	Amobarbital Carbamazepine Eslicarbazine Oxcarbazepine Phenobarbital Phenytoin Primidone Rufinamide
	Zonisamide			Rufinamide	
Antihypertensives	Aliskiren Amlodipine Diltiazem Eplerenone Felodipine Irbesartan Isradipine	Eplerenone Felodipine Isradipine	Diltiazem	Aliskiren	Aliskiren Enalapril Irbesartan Isradipine Non-DHP CCB Olmesartan Telmisartan Valsartan
				Enalapril Eplerenone Irbesartan Isradipine Non-DHP CCB Olmesartan Telmisartan	
Antipsychotics – first generation	Pimozide	Pimozide		Pimozide	
				Droperidol Thioridazine	
Antipsychotics – second generation	Paliperidone	Aripiprazole Quetiapine		Aripiprazole Clozapine Paliperidone Quetiapine	Paliperidone
Antiretrovirals	See HIV/HCV Coinfection Section				
Azole antifungals		Ketoconazole		Ketoconazole Posaconazole	
Benzodiazepines	Midazolam	Midazolam			

Bronchodilators				Theophylline	
Buprenorphine/naloxone					
Calcineurin inhibitors		Cyclosporine		Cyclosporine	Cyclosporine
		Tacrolimus		Tacrolimus	Tacrolimus
Cholesterol-lowering agents	Rosuvastatin	Atorvastatin Fluvastatin Gemfibrozil	Atorvastatin Fluvastatin Lovastatin	Atorvastatin Lovastatin Simvastatin	Rosuvastatin Pitavastatin
	Atorvastatin Fluvastatin Lovastatin Pitavastatin Pravastatin Simvastatin	Lovastatin Rosuvastatin Simvastatin	Pitavastatin Rosuvastatin Simvastatin	Ezetimibe Fluvastatin Gemfibrozil Pitavastatin Pravastatin Rosuvastatin	Pravastatin Atorvastatin Fluvastatin Lovastatin Simvastatin
Cisapride					
Ergot derivatives					
Ethinyl estradiol containing products					
Glucocorticoids		Dexamethasone		Dexamethasone	Dexamethasone
Heart failure agents		Bosentan	Bosentan	Bosentan	Bosentan
		Ambrisentan		Ambrisentan	Ambrisentan
Herbals	St. John's wort	St. John's wort	St. John's wort	St. John's wort	St. John's wort
Loop diuretics					
Macrolide antimicrobials	Telithromycin	Telithromycin		Erythromycin Telithromycin	Erythromycin Telithromycin
Phosphodiesterase -5 inhibitors					
Rifamycin antimicrobials	Rifabutin Rifampicin Rifapentine	Rifabutin Rifampicin Rifapentine	Rifabutin Rifampicin Rifapentine	Rifabutin Rifampicin Rifapentine	Rifabutin Rifampicin Rifapentine
		Rifaximin		Rifaximin	Rifaximin

H2RA=Histamine H₂ Antagonist; PPI=proton pump inhibitor; DHP CCB=dihydropyridine calcium channel blocker; Non-DHP CCB=non dihydropyridine calcium channel blocker

Green indicates coadministration is safe; **yellow** indicates a dose change or additional monitoring is warranted; and **red** indicates the combination should be avoided. Specific concomitant medications or medication classes with actual or theoretical potential for interaction are listed in the box.

The education of patients and caregivers about potential adverse effects of DAA therapy and their management is an integral component of treatment and is important for a successful outcome in all patient populations. During DAA treatment, individuals should be followed at clinically appropriate intervals to ensure medication adherence, assess adverse events and potential drug-drug interactions, and monitor blood test results necessary for patient safety. This includes on-treatment and post-treatment monitoring for hypoglycemia or subtherapeutic INR levels among patients taking diabetes medicines or warfarin, respectively. Real-world data indicate an association between DAA therapy and related changes in hepatic function and alterations in dose-response relationships with these medications ([Drazilova, 2018](#));

([Abdel Alem, 2017](#)); ([Rindone, 2017](#)); ([Pavone, 2016](#)); ([DeCarolis, 2016](#)); ([Soriano, 2016](#)). Inform patients on these medications about the potential for these developments; make dose adjustments as needed. The frequency and type of contact (eg, clinic visit, phone call, etc) are variable but need to be sufficient to assess patient safety and response to treatment, as outlined above.

Routine testing for HCV RNA during treatment is not recommended unless the ALT level fails to decline (when elevated) or there are concerns regarding patient adherence with DAA treatment. There are no data to support stopping treatment based on detectable HCV RNA during the first 4 weeks of treatment, or that detectable HCV RNA at this time point signifies medication nonadherence.

It is essential to test for HCV RNA 12 weeks (or longer) after treatment completion. Undetectable or unquantifiable HCV RNA 12 weeks or longer after treatment completion is defined as a sustained virologic response (SVR), which is consistent with cure of HCV infection. Virologic relapse is rare 12 weeks or longer after treatment completion ([Sarrazin, 2017](#)); ([Simmons, 2016](#)). Nevertheless, repeat quantitative HCV RNA testing can be considered at 24 or more weeks after completing treatment for patients in whom ALT increases to above the upper limit of normal.

During clinical trials with elbasvir/grazoprevir, with or without ribavirin, 1% of participants experienced ALT elevations from normal levels to >5 times the upper limit of normal, generally at or after treatment week 8. ALT elevations were typically asymptomatic and most resolved with ongoing therapy or completion of therapy. Higher rates of late ALT elevations occurred in females, those of Asian descent, and patients aged ≥ 65 years. Hepatic laboratory testing should be performed prior to therapy, at treatment week 8, and as clinically indicated. For patients receiving 16 weeks of therapy, additional hepatic laboratory testing should be performed at treatment week 12 ([Zepatier package insert, 2019](#)).

Patients being treated with amiodarone should not receive sofosbuvir-based regimens due to risk of life-threatening arrhythmias. Because of its long half-life, it is advised that persons should be off amiodarone for at least 6 months before initiating sofosbuvir. If the decision is made to start sofosbuvir in this setting, continued vigilance for bradycardia should be exercised.

Pregnancy and Nursing Mothers

Few adequate and well-controlled human studies are available to establish whether DAAs pose a risk to pregnancy outcomes or whether DAAs and their metabolites are present in breastmilk. An open-label, phase 1 study of HIV-negative pregnant women with chronic genotype 1 infection evaluated a 12-week course of ledipasvir/sofosbuvir initiated between 23 to 24 weeks of gestation ([Chappell, 2019](#)). Among 7 evaluable patients, all achieved SVR12 and adverse events related ledipasvir/sofosbuvir were \leq grade 2. All 7 participants delivered at term with undetectable HCV viral loads at delivery. One-year follow-up of the infants is ongoing.

Given the dearth of data on this topic, clinicians should discuss with female patients that DAAs should be used during pregnancy only if the potential benefit of DAA therapy justifies the potential risk of harm to the fetus. The health benefits of DAA therapy for nursing mothers should be weighed against the health benefits of breast feeding and the possible adverse effects of the DAA regimen on the breastfed child. Given the relatively short duration of treatment and the availability of ribavirin-free regimens in most patients, the potential risk of harms and benefits of delaying pregnancy until HCV DAA therapy is completed should be considered. For additional information about HCV and pregnancy, [click here](#).


Reactivation of HBV

Cases of HBV reactivation, occasionally fulminant, during or after DAA therapy have been reported in HBV/HCV coinfecting patients who were not receiving HBV suppressive therapy ([Mücke, 2018](#)); ([Bersoff-Matcha, 2017](#)); ([Chen, 2017](#)). Therefore, all patients initiating HCV DAA therapy should be assessed for HBV coinfection with HBsAg testing, and for evidence of prior infection with anti-HBc and anti-HBs testing. HBV vaccination is recommended for all susceptible individuals. Testing for HBV DNA should be performed prior to DAA therapy in patients who are HBsAg positive. HBsAg positivity does not represent a contraindication to HCV DAA therapy. Patients meeting criteria for treatment of active HBV infection should be started on HBV therapy at the same time or before HCV DAA therapy is initiated ([Terrault, 2018](#)).


Patients with a low or undetectable HBV DNA level can either receive prophylactic HBV treatment for the duration of DAA treatment until assessment for SVR12 or be monitored at regular intervals (usually not more frequently than every 4 weeks) for HBV reactivation with HBV DNA testing. If monitoring is elected, HBV treatment should be started if the HBV DNA level increases >10-fold or is >1000 IU/mL in a patient with undetectable or unquantifiable HBV DNA prior to DAA treatment. There are insufficient data to provide clear recommendations for the monitoring of HBV DNA among patients testing positive either for anti-HBc alone (isolated anti-HBc) or for anti-HBc and anti-HBs (resolved infection). However, the possibility of HBV reactivation should be considered in these patients in the event of an unexplained increase in liver aminotransferase levels during and/or after completion of DAA therapy.

Post-Treatment Follow-Up for Patients in Whom Treatment Failed

Recommended Monitoring for Patients in Whom Treatment Failed to Achieve a Sustained Virologic Response

RECOMMENDED	RATING 
Retreatment for chronic HCV is recommended utilizing the regimens recommended in the Retreatment section.	I, C
Disease progression assessment every 6 to 12 months with a hepatic function panel, complete blood count (CBC), and international normalized ratio (INR) is recommended.	I, C
Surveillance for hepatocellular carcinoma with liver ultrasound examination, with or without alpha fetoprotein (AFP), every 6 months is recommended for patients with cirrhosis ^a in accordance with the AASLD guidance on the diagnosis, staging, and management of hepatocellular carcinoma .	Low, Conditional ^b
For patients with cirrhosis, endoscopic surveillance for varices should be performed in accordance with the AASLD guidance on portal hypertension bleeding in cirrhosis .	Guidance ^b
<p>^a For decompensated cirrhosis, please refer to the appropriate section.</p> <p>^b Unlike the AASLD/IDSA HCV guidance, the AASLD guidelines for treatment of chronic hepatitis B uses the GRADE approach to rate recommendations; please see that document for further information about this rating system.</p>	


The Following Monitoring Is Not Recommended During or After Therapy

NOT RECOMMENDED	RATING 
Monitoring for HCV drug resistance-associated substitutions (RASs) during or after therapy is not recommended unless retreatment will be performed and RAS testing is recommended in advance of this therapy. See the Retreatment section for recommendations regarding RAS testing prior to retreatment. Additional information about RAS testing can be found in the HCV Resistance Primer .	IIb, C

Patients who do not achieve SVR retain the possibility of continued liver injury, progression of hepatic fibrosis, and the potential to transmit HCV infection to others. Such patients should be considered for retreatment per the [Retreatment of Persons in Whom Prior Therapy Has Failed](#) section.

Given that persons in whom treatment failed remain at risk for ongoing liver injury and liver fibrosis progression ([Dienstag, 2011](#)), these patients should be monitored for signs and symptoms of cirrhosis. Patients in whom antiviral therapy failed may harbor viruses that are resistant to one or more of the antivirals at the time of virologic breakthrough ([Lawitz, 2014a](#)); ([Schneider, 2014](#)). There is no evidence to date, however, that the presence of RASs results in more progressive liver injury than would have occurred if the patient did not have resistant viruses. Additional information about RASs and RAS testing can be found in the [HCV Resistance Primer](#) section. If there remains uncertainty regarding the applicability of RAS testing, consultation with an expert regarding the treatment of HCV infection may be useful.

Post-Treatment Follow-Up for Patients Who Achieved a Sustained Virologic Response

Recommended Follow-Up for Patients Who Achieved a Sustained Virologic Response (SVR)	
RECOMMENDED	RATING 
For noncirrhotic patients, recommended follow-up is the same as if they were never infected with HCV.	I, B
Assessment for HCV recurrence is recommended only if the patient develops unexplained hepatic dysfunction, or annual assessment if the patient has ongoing risk factors for HCV infection. In such cases, a quantitative HCV RNA test, rather than an HCV antibody test, is recommended to assess for HCV recurrence.	I, A
Surveillance for hepatocellular carcinoma is recommended for patients with cirrhosis ^a , in accordance with the AASLD guidance on the diagnosis, staging, and management of hepatocellular carcinoma .	Strong, Moderate ^b
For cirrhotic patients, upper endoscopic surveillance is recommended in accordance with the AASLD guidance on portal hypertension bleeding in cirrhosis .	Guidance ^b
Assessment for other causes of liver disease is recommended for patients who develop persistently abnormal liver tests after achieving SVR.	I, C
^a For decompensated cirrhosis , please refer to the appropriate section. ^b Unlike the AASLD/IDSA HCV guidance, the AASLD guidelines for treatment of chronic hepatitis B uses the GRADE approach to rate recommendations; please see that document for further information about this rating system.	

Patients with undetectable serum HCV RNA, as assessed by a sensitive polymerase chain reaction (PCR) assay, ≥ 12 weeks after treatment completion are deemed to have achieved SVR (cure). The likelihood of achieving SVR with DAA therapy among adherent, immunologically competent, treatment-naïve patients with compensated liver disease generally exceeds 95%. Among patients who achieved SVR with peginterferon/ribavirin treatment, more than 99% have remained free of HCV infection when followed for 5 years after treatment completion ([Manns, 2013](#)). Thus, achieving SVR is considered a virologic cure of HCV infection. SVR typically aborts progression of liver injury with regression of liver fibrosis in most (but not all) treated patients ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)). Liver fibrosis and liver function test results improve in most patients who achieve SVR ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)). Because of lack of progression, noncirrhotic patients who achieve SVR should receive standard medical care that is recommended for patients who were never infected with HCV unless they

remain at risk for non-HCV-related liver disease, such as nonalcoholic fatty liver disease or alcoholic liver disease.

Among cirrhotic patients who achieve SVR, decompensated liver disease (with the exception of hepatocellular carcinoma [HCC]) rarely develops during follow-up and overall survival is prolonged ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)). Bleeding from esophageal varices is rare after SVR ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)). Cirrhotic patients should undergo surveillance endoscopy every 2 years if known to have small varices and every 3 years in the absence of known varices in accordance with AASLD guidance on portal hypertension bleeding ([Garcia-Tsao, 2017](#)).


Importantly, cirrhotic patients remain at risk for developing HCC and should, therefore, undergo surveillance for HCC every 6 months utilizing ultrasound (with or without AFP testing) despite the lowered risk that results after viral eradication ([Marrero, 2018](#)). Although multiple studies of cirrhotic patients who achieved SVR with peginterferon/ribavirin reported a reduction in the risk of developing HCC ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)) and a meta-analysis of persons achieving SVR with DAAs found that the HCC risk did not exceed that seen in patients who experienced SVR with interferon-based treatment after adjustment for baseline risk factors for HCC ([Waziry, 2017b](#)), one report found a higher than expected frequency of HCC in patients with HCV-related cirrhosis despite successful DAA treatment ([Reig, 2016](#)). However, a prospective observational study of 3045 cirrhotic patients found an adjusted hazard ratio for HCC of 0.57 (95% CI 0.40 to 0.81) following DAA-based therapy, implying a 43% reduction in HCC incidence ([Carrat, 2019](#)).

Bleeding from esophageal varices is uncommon after SVR ([Morgan, 2013](#)); ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([Singal, 2010](#)); ([George, 2009](#)). Nevertheless, patients with compensated cirrhosis who achieve SVR should continue to receive endoscopic surveillance for esophageal varices, in accordance with the AASLD guidance on portal hypertension bleeding ([Garcia-Tsao, 2017](#)). Current AASLD recommendations for patients with compensated cirrhosis without known varices is surveillance endoscopy every 2 years if there is evidence of ongoing liver injury from associated conditions, such as obesity or alcohol use, and every 3 years if liver injury is quiescent, such as after alcohol abstinence. Patients with compensated cirrhosis and known varices should undergo surveillance endoscopy annually if there is evidence of ongoing liver injury from associated conditions, such as obesity or alcohol use, and every 2 years if liver injury is quiescent, such as after alcohol abstinence.

Patients in whom SVR is achieved but who have another potential cause of liver disease (eg, excessive alcohol use, metabolic syndrome with or without proven fatty liver disease, or iron overload) remain at risk for hepatic fibrosis progression. It is recommended that such patients be educated about the risk of liver disease and monitored for liver disease progression with periodic physical examination, blood tests, and potentially, tests for liver fibrosis by a liver disease specialist.

Patients who achieve SVR can have HCV recurrence due to reinfection or late relapse ([Sarrazin, 2017](#)); ([Simmons, 2016](#)). A systematic review suggests 5-year recurrence risks of 1%, 11%, and 15% in monoinfected low-risk HCV, monoinfected high-risk HCV (ie, people who currently or formerly injected drugs, imprisonment, or men who have sex with men [MSM]), and HIV/HCV coinfecting patients, respectively ([Simmons, 2016](#)). At least annual testing for HCV reinfection among patients with ongoing risk for HCV infection (eg, injection drug use or high-risk sexual exposure) is recommended. A flare in liver enzyme levels should prompt immediate evaluation for HCV reinfection (see [Management of Acute HCV Infection](#)). Because HCV antibody remains positive in most patients following SVR, testing for HCV recurrence using an assay that detects HCV RNA (ie, a quantitative HCV-RNA test) is recommended.

Monitoring for HCV During Chemotherapy and Immunosuppression


NOT RECOMMENDED	RATING 
Prospective monitoring for HCV recurrence among patients who achieved SVR and are receiving immunosuppressive drug therapy (eg, systemic corticosteroids, antimetabolites, chemotherapy, biologics agents, etc) is not routinely recommended.	III, C

Monitoring for HCV During Chemotherapy and Immunosuppression


Acute liver injury is common among patients receiving chemotherapy or immunosuppressive agents. Testing for hepatitis viruses should be included in the laboratory assessment of the cause of acute liver injury in these patients. Approximately 23% of patients with active HCV infection—especially those with a hematologic malignancy—experience a flare in their HCV RNA level (>10-fold) during chemotherapy. An ALT level increase is less common and clinical symptoms of hepatitis are uncommon ([Torres, 2018](#)). Among patients who have recovered from HCV infection, either spontaneously or with DAA treatment, reactivation of HCV (ie, detectable HCV RNA) during chemotherapy is distinctly uncommon and is not anticipated to occur since there is no residual reservoir for the virus. Thus, in this latter group, routine testing for HCV RNA during immunosuppressive treatment or prophylactic administration of antivirals during immunosuppressive treatment is not recommended.

Additional Considerations If Treatment Includes Ribavirin

Recommended Monitoring During Antiviral Therapy That Includes Ribavirin

RECOMMENDED	RATING 
More frequent assessment for drug-related adverse effects (ie, CBC for patients receiving ribavirin) is recommended as clinically indicated.	I, C

Recommended Monitoring for Pregnancy-Related Issues Prior to and During Antiviral Therapy That Includes Ribavirin

RECOMMENDED	RATING 
Women of childbearing potential and their partners should not receive ribavirin during or for at least 6 months prior to pregnancy.	I, C
Women of childbearing potential should be counseled not to become pregnant while receiving a ribavirin-containing antiviral regimen, and for at least 6 months after stopping the regimen.	I, C
Male partners of women of childbearing potential should be cautioned to prevent pregnancy while they are receiving a ribavirin-containing antiviral regimen, and for up to 6 months after stopping the regimen.	I, C
Serum pregnancy testing is recommended for women of childbearing potential prior to beginning treatment with a regimen that includes ribavirin.	I, C
Assessment of contraceptive use and of possible pregnancy is recommended at appropriate intervals during (and for 6 months after) ribavirin treatment for women of childbearing potential, and for female partners of men who receive ribavirin treatment.	I, C

Recommended Monitoring for Pregnancy-Related Issues Prior to and During Antiviral Therapy That Includes Ribavirin

Ribavirin causes hemolysis. Patients receiving ribavirin should have hemoglobin levels checked during treatment, often after 2 weeks, and the ribavirin dose reduced if the patient develops significant anemia, often defined as hemoglobin <10 g/dL.

Ribavirin causes fetal death and fetal abnormalities in animals. Ribavirin should not be administered to pregnant women or to women who might become pregnant during or for 6 months after completing ribavirin treatment. Similarly, ribavirin may cause birth defects in offspring of women whose partner was receiving ribavirin when the woman became pregnant. In the very rare instances when ribavirin is used, it is imperative for persons of childbearing potential to use at least 2 reliable forms of effective contraception during treatment and for a period of 6 months thereafter. It is recommended that the healthcare practitioner document the discussion of the potential teratogenic effects of ribavirin in the patient's medical record.

Last update: August 27, 2020

HCV Resistance Primer

Introduction

Understanding principles of the emergence of drug-resistant viruses is critical when using targeted antiviral therapies. The best example of these principles can be gleaned from the study of HIV. Like HIV, HCV is an approximately 9.5 kilobase RNA virus that replicates very rapidly (billions of viruses daily). The production of each new virus is performed by an enzyme that results in 1 to 3 errors per replication cycle, on average. Many of these errors either have no effect on the progeny virus product or result in progeny viruses that are nonreplication competent (ie, dead viruses). For some newly produced viruses, however, the transcription errors result in changes in critical coding regions that may, by chance, change the susceptibility of the virus to 1 or more drugs used to treat the virus. The emergence of such drug-resistant viruses most often occurs when drug levels are subtherapeutic, thereby creating selective pressure for the resistant viruses to emerge as the dominant species. These newly formed resistant viruses have a selective growth advantage that allows them to replicate in the presence of antiviral drugs. In a subset of patients with chronic HCV infection, viral variants harboring substitutions associated with resistance to HCV directing-acting antivirals (DAAs) are detectable prior to antiviral therapy and, particularly in the case of NS5A inhibitor-containing regimens, may negatively impact treatment response. These substitutions often are referred to as baseline resistance-associated substitutions (RASs).

In the case of HCV DAAs, resistant viruses are also selected for and/or enriched in patients for whom a DAA regimen fails. These viruses contain substitutions that are designated as treatment-emergent (or treatment-selected) RASs. NS5A and NS3 RASs are frequently selected in patients with failure of NS5A or NS3 inhibitor-containing regimens, respectively. In contrast, NS5B nucleotide RASs are rarely detected (1% of failures) even after exposure to a failing DAA regimen containing a nucleotide inhibitor ([Svarovskaia, 2014](#)); ([Wyles, 2018b](#)). This is likely due to the highly conserved catalytic site region that nucleotides bind, making substitutions in this region extremely rare—often referred to as a high barrier to resistance. Additionally, any such substitution would likely render the virus replication incompetent. Compounding the clinical impact of NS5A RASs is their ability to maintain high replication competence (aka, relative fitness) in the absence of continued drug pressure, allowing them to remain the dominant viral quasispecies for prolonged periods (years) relative to NS3 protease or NS5B nucleotide polymerase inhibitor RASs, which are typically less fit and tend to disappear over

several months, being overcome by more fit wild-type virus species.

The magnitude of the negative impact of both baseline and selected RASs on treatment outcome varies according to regimen (ie, coadministered drugs); patient factors that impact treatment response (eg, cirrhosis); and the fold change decrease in potency conferred by the specific RAS(s). Given these considerations, RAS testing alone will not dictate optimal DAA regimen selection. In addition, a drug predicted to suffer a significant loss of potency in the presence of a RAS still may be used in specific clinical settings/regimens.

Terminology, Thresholds of Clinical Relevance, and Assays

Terminology

1. Naming Convention for Hepatitis C Proteins

The hepatitis C genome codes for approximately 5 HCV-specific proteins, which are essential to: 1) form the viral structure (core and envelope proteins); 2) cut the HCV polyprotein; 3) provide enzymatic functions for replication and escape from the innate immune response (NS3/NS4A protease); 4) replicate the HCV RNA (NS5B RNA-dependent RNA polymerase); and 5) bind the HCV replication complex during replication and assembly (NS5A).

2. Polymorphism (Substitution)

A reference (or consensus) nucleotide—and therefore amino acid sequence—has been defined for each HCV genotype. A polymorphism (or substitution) is a difference in an amino acid at a defined position of the HCV protein between a patient's HCV and the reference HCV protein. Substitution is the preferred terminology among most experts. However, the US Food and Drug Administration currently uses the term polymorphism.

To define a polymorphism, it is necessary to define: the HCV genotype (eg, genotype 1, 2, 3, etc) and subtype (eg, 1a vs 1b); the HCV protein (eg, NS5A); and the amino acid position (eg, 93). Polymorphisms are reported as letter-number-letter (eg, Y93H). The first letter refers to the amino acid typically expected for that position in the reference protein. The number refers to the amino acid position, and the final letter refers to the amino acid that is found in the patient's HCV isolate. Thus, NS5A Y93H refers to amino acid position 93 of the NS5A protein. The amino acid at this position in the reference strain is Y (ie, tyrosine) and the amino acid in the tested strain is H (ie, histidine). For some patients, multiple variants are present and several amino acids may be found at a given position. Thus, it is possible to have a virus with NS5A Y93H/M. Such a patient would have viruses with the amino acids histidine (H) or methionine (M) at position 93 of the NS5A protein.

3. Resistance-Associated Substitutions

A resistance-associated substitution describes any amino acid change from the consensus sequence at a position that has been associated with reduced susceptibility of a virus to 1 or more antiviral drugs. A specific RAS may or may not confer a phenotypic loss of susceptibility to other/multiple antiviral agents.

4. Drug-Class RASs

Drug-class RASs are amino acid substitutions that reduce the susceptibility of a virus to any (and at least 1) member of a drug class or, alternatively, the viral variants with reduced susceptibility that carry these substitutions. Class RASs may or may not confer resistance to a specific drug in that class.

5. Drug-Specific RASs

Drug-specific RASs are amino acid substitutions that reduce the susceptibility of a virus to a specific drug. When assessing the potential clinical impact of RASs on a given regimen, drug-specific RASs should be used. In an HCV-infected population not previously exposed to a DAA drug or class, drug-specific RASs will be found less frequently than class RASs.

Thresholds of Clinical Relevance

HCV resistance to DAAs is a rapidly evolving field with demonstrated clinical impact in specific situations with currently available DAA regimens. Presently, the most clinically significant RASs are in the NS5A position for genotypes 1a and 3.

Data from clinical trials have demonstrated that RASs are commonly, but not always, found at the time of virologic failure. Viruses that are resistant to NS3/4A protease inhibitors seem to be less fit and may disappear from peripheral blood within a few weeks to months, whereas NS5A inhibitor-resistant viruses may persist for years, which could have implications for treatment and retreatment.

In general, drug-specific RASs need to be present in at least 15% of the viruses of a given patient to reduce the likelihood of achieving SVR ([Pawlotsky, 2016](#)). Drug-specific RASs that are found at a lower frequency may not convey sufficient resistance to reduce SVR with currently available DAA regimens.

Assays

Methods to detect RASs include population sequencing (aka, Sanger sequencing) and deep sequencing (aka, next generation sequencing [NGS]). Both methods depend on sequencing the HCV RNA, calculating the amino acid sequence, and then inferring the presence of RASs. The methods differ in their sensitivity for detecting RASs. For the purposes of clinical care and decisions regarding which DAA regimen to use, both methods can be considered equivalent if a $\geq 15\%$ cut point is used for determination of RASs by NGS. Recent studies have shown that NGS at a 1% level of sensitivity often result in the identification of additional RASs that are not associated with clinical failure ([Jacobson, 2015b](#)); ([Sarrazin, 2016](#)); ([Zeuzem, 2017](#)).

1. Genotypic Analysis

a. Population-Based Sequencing (Sanger)

Population sequencing of the HCV coding region of interest may be performed using reverse transcription polymerase chain reaction (PCR) and standard Sanger sequencing of the bulk PCR product. The sensitivity for detection of resistance substitutions varies but is generally 15% to 25%. As a standard, substitutions are reported as differences compared with a genotype-specific, wild-type strain.

b. Deep Sequencing Analysis

NGS (deep sequencing approaches) can increase the sensitivity of detection for minor variants. After sequencing HCV coding regions using PCR, a software algorithm is used to process and align sequencing data via a multistep method to identify the substitutions present at a predetermined level. This level, or threshold, can vary but is often set as low as $>1\%$ for research purposes. To approximate results obtained by population sequencing, NGS thresholds are often set to $\geq 10\%$.

2. Phenotypic Analysis


Phenotypic analysis involves laboratory techniques whereby the degree of drug resistance conferred by an amino acid change as well as the replicative capacity (fitness) of a particular RAS can be estimated in the presence of a wild-type or consensus strain. These research techniques are not routinely used for clinical practice. To assess the

level of resistance, RASs are typically introduced as point mutations into the backbone of an existing standard HCV genome within an existing cell culture/replicon or enzyme-based assay. Isolates harboring these RASs are then challenged by appropriate antiviral agents at increasing concentrations and fold changes—based on EC₅₀ or IC₅₀ and EC₉₀ or IC₉₀ values—are determined for inhibition of replication or enzyme activity, respectively, in comparison to wild-type virus. Comparison of replication levels for variants and wild-type constructs in the absence of drug allows for estimation of fitness.

3. Assay Summary Points

- Either population sequencing or deep sequencing can be used to detect the presence of RASs in NS3, NS5A, and NS5B.
- For clinical decisions, population sequencing or deep sequencing with at least 15% prevalence of RASs as the cutoff is recommended. The presence of RASs with <15% prevalence should not be considered clinically significant.
- When assessing the potential clinical effect of RASs, it is important to determine the drug-specific RASs.

Resistance Testing in Clinical Practice

Regimen-Specific Recommendations for Use of RAS Testing in Clinical Practice	
RECOMMENDED	RATING 
<p>Elbasvir/grazoprevir NS5A RAS testing is recommended for genotype 1a-infected, treatment-naïve or -experienced patients being considered for elbasvir/grazoprevir. If present, a different regimen should be considered.</p>	I, A
<p>Ledipasvir/sofosbuvir NS5A RAS testing can be considered for genotype 1a-infected, treatment-experienced patients with and without cirrhosis being considered for ledipasvir/sofosbuvir. If clinically important^a resistance is present, a different recommended therapy should be used.</p>	I, A
<p>Sofosbuvir/velpatasvir NS5A RAS testing is recommended for genotype 3-infected, treatment-naïve patients with cirrhosis and treatment-experienced patients (without cirrhosis) being considered for 12 weeks of sofosbuvir/velpatasvir. If Y93H is present, weight-based ribavirin should be added or another recommended regimen should be used.</p>	I, A
<p>^a Clinically important = ≥100-fold shift in the in vitro EC₅₀ to ledipasvir</p>	

Resistance testing is most important in clinical practice when the results would modify treatment management by impacting the duration of therapy and/or inclusion of ribavirin, or result in selection of alternative therapy. Unfortunately the

utility of RAS testing at this time varies by both patient characteristics and DAA regimen.

Approaches to Overcome Resistance

Data for currently approved DAAs provide limited insight on optimal retreatment approaches for patients with a previous DAA therapy failure and high fold change RASs, particularly those in NS5A. Until regimens combining multiple drugs predicted to be active (based on the available resistance profile) are available and adequate phase 2/3 studies in DAA treatment failure populations are accomplished, other aspects of therapy must be optimized in treatment-experienced patients with RASs. In general, optimization involves appropriately characterizing the patient along with use of an extended duration of therapy and the addition of ribavirin (unless an absolute contraindication to ribavirin exists).

Characterizing Patients at Risk

The characteristics that increase the risk of DAA treatment failure are different for each oral regimen. Thus, understanding the population at risk is imperative. Generally, this requires accurate assessment of liver fibrosis and clarification of prior therapy.

Virus

Determination of HCV genotype, subtype, and baseline RASs may be necessary to fully characterize a patient's risk for therapeutic failure and optimize the treatment approach.

Treatment Duration

The duration of therapy should always be optimized to attain a cure. Although short-duration therapy has been associated with a higher chance of relapse, careful selection of patients for shortened therapy may minimize relapse risk and lead to significant cost savings. In contrast, extension of therapy (often to 24 weeks) in conjunction with the addition of ribavirin has been associated with reasonable SVR rates during retreatment of patients with past DAA therapy failure, even in the presence of significant drug-specific RASs prior to retreatment ([Cooper, 2016](#)); ([Gane, 2017](#)).

Ribavirin

The addition of ribavirin increases SVR in patient populations with an increased risk for treatment failure (eg, decompensated cirrhosis). It also improves SVR rates among patients with baseline NS5A RASs and prior DAA treatment failure.

Complementary Therapy

Although data are limited, patients with multiclass RASs can achieve SVR by combining triple or quadruple drug class regimens (see section on [retreatment](#) in prior DAA failure). This approach may become less necessary with the approval of standalone dual- or triple-drug regimens composed of second-generation protease and NS5A inhibitors with improved activity against common RASs.

Considerations With Current Antiviral Regimens

Elbasvir/Grazoprevir

Elbasvir/grazoprevir is indicated for treatment-naïve and -experienced patients with genotype 1 or 4. The presence of NS3 RASs has no significant impact on SVR12 in patients treated with elbasvir/grazoprevir. The presence of NS5A RASs has no significant impact in genotype 1b infection.

In treatment-naive, genotype 1a patients (with or without cirrhosis) treated with 12 weeks of therapy, the presence of NS3 RASs has no impact ([Zeuzem, 2015](#)). In treatment-naive or prior relapse patients treated for 12 weeks with elbasvir/grazoprevir without ribavirin, the presence of high fold change NS5A RASs (at amino acid positions 28, 30, 31, and 93) decreased SVR to 58% (14/24) compared to 98% SVR in those without NS5A RASs. The presence of NS5A RASs had a similar impact on treatment-experienced patients (with or without cirrhosis) who received 12 weeks of elbasvir/grazoprevir without ribavirin (SVR12 29% vs 97%, respectively) ([Jacobson, 2015b](#)).

Glecaprevir/Pibrentasvir

In a study of the resistance profiles of glecaprevir and pibrentasvir using cell cultures ([Ng, 2017](#)), selection of genotypes 1a, 1b, 2a, 3a, 4a, and 6a replicons for reduced susceptibility to glecaprevir resulted in the emergence of RASs at A156 or D/Q168. The A156 RAS resulted in the greatest reductions (>100-fold) in glecaprevir susceptibility. The D/Q168 RAS had varying effects on glecaprevir susceptibility depending on genotype/subtype and specific amino acid change. The greatest reductions (>30-fold) were observed in genotypes 1a (D168F/Y), 3a (Q168R), and 6a (D168A/G/H/V/Y). These RASs, however, are rarely detected clinically. Pibrentasvir selected no viable colonies in genotype 1b, 2b, 4a, 5a, and 6a. Of the few RASs selected by pibrentasvir, Y93H/N conferred <7-fold resistance.

The presence of baseline RASs had minimal impact on SVR rates with glecaprevir/pibrentasvir in registration trials that predominantly enrolled noncirrhotic patients. In a pooled analysis of NS3/4A protease inhibitor- and NS5A inhibitor-naive patients who received glecaprevir/pibrentasvir in phase 2 and 3 studies ([Forns, 2017](#)); ([Foster, 2017](#)); ([Asselah, 2018b](#)); ([Zeuzem, 2016](#)); ([Kwo, 2017b](#)), baseline RASs in patients with genotype 1, 2, 4, 5, or 6 infection had no impact on SVR12 ([Krishnan, 2018](#)). Among treatment-naive genotype 3 patients without cirrhosis who received glecaprevir/pibrentasvir for 8 weeks, the A30K polymorphism was detected in 10%, of whom 78% achieved SVR12. There are insufficient data to characterize the impact of A30K in genotype 3 patients with cirrhosis or prior treatment experience. All genotype 3 patients with Y93H prior to treatment achieved SVR12.

Ledipasvir/Sofosbuvir

Several comprehensive analyses of genotype 1 patients treated with ledipasvir/sofosbuvir in phase 2 and phase 3 studies have helped clarify the impact of baseline RASs on SVR rates with this regimen ([Sarrazin, 2016](#)); ([Zeuzem, 2017](#)). In a pooled analysis of patients with genotype 1a or 1b who received ledipasvir/sofosbuvir, 93.5% (316/338) of those with baseline NS5A RASs achieved SVR12 compared to an SVR12 of 98.4% (1,741/1,770) in patients without baseline NS5A RASs ([Sarrazin, 2016](#)). In this analysis, the reduction in SVR was driven predominantly by patients with genotype 1a NS5A RASs. The SVR12 rates for genotype 1a patients with and without NS5A RASs were 92.3% and 98.3%, respectively. A slightly lower SVR12 of 90% was observed for genotype 1a patients with NS5A RASs using a 15% deep sequencing cutoff value.

Notably, other factors further delineated populations at risk for relapse in this analysis, including high-level baseline NS5A RASs (>100-fold resistance with Q30H/R, L31M/V, and Y93C/H/N in genotype 1a) and a shorter duration therapy (8 weeks or 12 weeks vs 24 weeks). SVR12 rates were 97.4% to 100% in treatment-experienced patients without NS5A RASs or with RASs with <100-fold resistance treated with ledipasvir/sofosbuvir for 12 weeks or 24 weeks. When RASs with >100-fold resistance were present, however, SVR12 dropped to 64.7% (11/17) with 12 weeks of therapy compared to 100% (6/6) with 24 weeks of therapy. In this small subset of patients, the addition of ribavirin did not appear to offer the same benefit as extension of therapy to 24 weeks in this pooled analysis. SVR12 was 81.8% in those with >100-fold NS5A resistance who received 12 weeks of ledipasvir/sofosbuvir with ribavirin. In contrast, in the SIRIUS trial, all 8 treatment-experienced cirrhotic patients with >100-fold resistance treated for 12 weeks with ledipasvir/sofosbuvir plus ribavirin achieved SVR12.

Sofosbuvir/Velpatasvir

Sofosbuvir/velpatasvir is a pangenotypic therapy indicated for treatment-naive and -experienced patients with or without cirrhosis. In the ASTRAL studies, the presence of NS5A RASs had no impact on SVR12 for patients with genotype 1, 2, 4, 5, or 6 infection treated with 12 weeks of sofosbuvir/velpatasvir ([Hézode, 2018](#)). The presence of Y93H in genotype 3 patients decreased the SVR12 to 84% (21/25 patients) compared to 97% (242/249) in those without this RAS ([Foster, 2015a](#)). This appeared to be more impactful in patients with cirrhosis and/or prior treatment experience with an interferon-based regimen. Ribavirin was not used in these trials. However, a subsequent trial that randomized patients with genotype 3 and cirrhosis to sofosbuvir/velpatasvir with or without ribavirin demonstrated lower relapse rates in patients receiving ribavirin ([Esteban, 2018](#)).

Sofosbuvir/Velpatasvir/Voxilaprevir

Sofosbuvir/velpatasvir/voxilaprevir fills an important role as a pangenotypic regimen for patients who have experienced treatment failure with DAA therapy. Although data are limited, the presence of NS3, NS5A, or NS5B RASs prior to treatment did not influence the likelihood of SVR12, and 12 weeks of treatment produced a high SVR12 (96%) in DAA-experienced patients. RAS testing has not been demonstrated to impact SVR rates with sofosbuvir/velpatasvir/voxilaprevir therapy ([Bourlière, 2017](#)); ([Sarrazin, 2018](#)).

Table 1. Most Common, Clinically Important RASs by DAA, Genotype, and Fold Change

DAA	Genotype 1a				Genotype 1b		Genotype 3a	
	M28T	Q30R	L31M/V	Y93H/N	L31V/I	Y93H/N	A30K	Y93H
Ledipasvir	20x	>100x	>100x / >100x	>1000x / >10,000x	>100x >50x	>100x / --	NA	NA
Elbasvir	20x	>100x	>10x >100x	>1000x / >1000x	<10x	>100x / --	50x	>100x
Velpatasvir	<10x	<3x	20x / 50x	>100x / >1000x	<3x	<3x / --	50x	>100x
Pibrentasvir	<3x	<3x	<3x	<10x	<3x	<3x	<3x	<3x

Color Key: light green = <3-fold change; dark green = <10-fold change; orange = >10- to 100-fold change; pink = >100-fold change

Table 2. Clinically Important RASs by DAA Regimen and Genotype

DAA Regimen	Genotype		
	1a	1b	3
Ledipasvir/sofosbuvir	Q30H/R L31M/V Y93C/H/N	L31V ?Y93H	NA
Elbasvir/grazoprevir	M28A/T Q30H/R L31M/V Y93C/H/N	Y93H	NA
Sofosbuvir/velpatasvir	NA	NA	Y93H

Table 3. NS5A RAS Testing Recommendations Prior to Initiation of DAA Treatment Among Genotype 1 Patients by DAA Regimen, Virus Subtype, Prior Treatment Status, and Cirrhosis Status

DAA Regimen	1b TN ^a or TE ^b	1a TN	1a TE No Cirrhosis	1a TE Cirrhosis
Ledipasvir/sofosbuvir	No	No	Yes	Yes
Elbasvir/grazoprevir	No	Yes	Yes	Yes
Sofosbuvir/velpatasvir	No	No	No	No
^a TN = treatment naive ^b TE = treatment experienced				

Last update: November 6, 2019

Related References

[AASLD/IDSA HCV guidance panel, Recommendations for testing, managing, and treating hepatitis C website. Published](#)

[2017. Accessed June 13, 2017](#) [Internet]. 2017. <http://hcvguidelines.org/>

Abdel-Alem S, ElSharkawy AM, Fouad R, et al. [Improvement of glycemic state among responders to sofosbuvir-based treatment regimens: single center experience](#). J Med Virol. 2017;89:2181-2187.

Aberg JA, Gallant JE, Ghanem KG, Emmanuel P, Zingman BS, Horberg MA. [Infectious Disease Society of America. Primary care guidelines for the management of persons infected with HIV: 2013 update by the HIV Medicine Association of the Infectious Diseases Society of America](#). Clin Infect Dis. 2014;58(1):e1-34.

Afdhal NH, Bacon BR, Patel K, Lawitz EJ, Gordon SC, Nelson DR, et al. [Accuracy of fibroscan, compared with histology, in analysis of liver fibrosis in patients with hepatitis B or C: a United States multicenter study](#). Clin Gastroenterol Hepatol. 2015;13(4):772-779.

Agnello V, Chung RT, Kaplan LM. [A role for hepatitis C virus infection in type II cryoglobulinemia](#). N Engl J Med. 1992;327(21):1490-1495.

Allen SA, Spaulding AC, Osei AM, Taylor LE, Cabral AM, Rich JD. [Treatment of chronic hepatitis C in a state correctional facility](#). Ann Intern Med. 2003;138(3):187-190.

Allison RD, Tong X, Moorman AC, Ly KN, Rupp L, Xu F, et al. [Increased incidence of cancer and cancer-related mortality among persons with chronic hepatitis C infection, 2006-2010](#). J Hepatol. 2015;63(4):822-828.

Alter MJ, Kuhnert WL, Finelli L, Finelli L. [Guidelines for laboratory testing and result reporting of antibody to hepatitis C virus](#). MMWR Recomm Rep. 2003;52(RR-3):1-13, 15.

Amon JJ, Garfein RS, Ahdieh-Grant L, Armstrong GL, Ouellet LJ, Latka MH, et al. [Prevalence of hepatitis C virus infection among injection drug users in the United States, 1994-2004](#). Clin Infect Dis. 2008;46(12):1852-1858.

Arase Y, Suzuki F, Suzuki Y, Akuta N, Kobayashi M, Kawamura Y, et al. [Sustained virological response reduces incidence of onset of type 2 diabetes in chronic hepatitis C](#). Hepatology. 2009;49(3):739-744.

Arora S, Thornton K, Murata G, Deming P, Kalishman S, Dion D, et al. [Outcomes of treatment for hepatitis C virus infection by primary care providers](#). N Engl J Med. 2011;364(23):2199-2207.

Aspinall EJ, Corson S, Doyle JS, Grebely J, Hutchinson SJ, Dore GJ, et al. [Treatment of hepatitis C virus infection among people who are actively injecting drugs: a systematic review and meta-analysis](#). Clin Infect Dis. 2013;57(Suppl 2):S80-S89.

Asselah T, Kowdley KV, Zadeikis N, Wang S, Hassanein T, Horsmans Y, et al. [Efficacy of glecaprevir/pibrentasvir for 8 or 12 weeks in patients with hepatitis C virus genotype 2, 4, 5, or 6 infection without cirrhosis](#). Clin Gastroenterol Hepatol. 2018;16(3):417-426.

Assoumou SA, Tasillo A, Leff JA, Schackman BR, Drainoni M-L, C. Horsburgh R, et al. [Cost-effectiveness of one-time hepatitis C screening strategies among adolescents and young adults in primary care settings](#). Clin Infect Dis. 2018;66(3):376-384.

- Backus LI, Boothroyd DB, Phillips BR, Belperio P, Halloran J, Mole LA. [A sustained virologic response reduces risk of all-cause mortality in patients with hepatitis C](#). Clin Gastroenterol Hepatol. 2011;9(6):509-516.
- Barocas JA, Tasillo A, Eftekhari-Yazdi G, et al. [Population-level outcomes and cost-effectiveness of expanding the recommendation for age-based hepatitis C testing in the United States](#). Clin Infect Dis. 2018;67(4):549-556.
- Barua S, Greenwald R, Grebely J, Dore GJ, Swan T, Taylor LE. [Restrictions for Medicaid reimbursement of sofosbuvir for the treatment of hepatitis C virus infection in the United States](#). Ann Intern Med. 2015;163(3):215-223.
- Bedossa P, Dargère D, Paradis V. [Sampling variability of liver fibrosis in chronic hepatitis C](#). Hepatology. 2003;38(6):1449-1457.
- Bellentani S, Pozzato G, Saccoccio G, Crovatto M, Croce LS, Mazzoran L, et al. [Clinical course and risk factors of hepatitis C virus related liver disease in the general population: report from the Dionysos study](#). Gut. 1999;44(6):874-880.
- Benhamou Y, Bochet M, Di Martino V, Charlotte F, Azria F, Coutellier A, et al. [Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. The Multivirc group](#). Hepatology. 1999;30(4):1054-1058.
- Benhamou Y, Di Martino V, Bochet M, Colombet G, Thibault V, Liou A, et al. [Factors affecting liver fibrosis in human immunodeficiency virus and hepatitis C virus-coinfecting patients: impact of protease inhibitor therapy](#). Hepatology. 2001;34(2):283-287.
- Berenguer M, Palau A, Aguilera V, Rayon JM, Juan FS, Prieto M. [Clinical benefits of antiviral therapy in patients with recurrent hepatitis C following liver transplantation](#). Am J Transplant. 2008;8(3):679-687.
- Berenguer J, Álvarez-Pellicer J, Martín PM, López-Aldeguer J, Von-Wichmann MA, Quereda C, et al. [Sustained virological response to interferon plus ribavirin reduces liver-related complications and mortality in patients coinfecting with human immunodeficiency virus and hepatitis C virus](#). Hepatology. 2009;50(2):407-413.
- Berenguer M, Schuppan D. [Progression of liver fibrosis in post-transplant hepatitis C: mechanisms, assessment and treatment](#). J Hepatol. 2013;58(5):1028-1041.
- Bersoff-Matcha SJ, Cao K, Jason M, Ajao A, S. Jones C, Meyer T, et al. [Hepatitis B virus reactivation associated with direct-acting antiviral therapy for chronic hepatitis C virus: a review of cases reported to the US Food and Drug Administration adverse event reporting system](#). Ann Intern Med. 2017;166(11):792-798.
- Beste LA, Glorioso TJ, Ho PM. [Telemedicine specialty support promotes hepatitis C treatment by primary care providers in the Department of Veterans Affairs](#). Am J Med. 2017;130(4):432-438.e3.
- Bochud PY, Cai T, Overbeck K, Bochud M, Dufour JF, Mullhaupt B, et al. [Genotype 3 is associated with accelerated fibrosis progression in chronic hepatitis C](#). J Hepatol. 2009;51(4):655-666.
- Boerekamps A, van den Berk GE, Lauw FN, Leyten EM, van Kasteren ME, van Eeden A, et al. [Declining hepatitis C virus \(HCV\) incidence in Dutch human immunodeficiency virus-positive men who have sex with men after unrestricted access to HCV therapy](#). Clin Infect Dis. 2017;66(9):1360-1365.

- Bonder A, Afdhal NH. [Utilization of FibroScan in clinical practice](#). *Curr Gastroenterol Rep*. 2014;16(2):372.
- Bonkovsky HL, Snow KK, Malet PF, Back-Madruga C, Fontana RJ, Sterling RK, et al. [Health-related quality of life in patients with chronic hepatitis C and advanced fibrosis](#). *J Hepatol*. 2007;46(3):420-431.
- Boscarino JA, Lu M, Moorman AC, Gordon SC, Rupp LB, Spradling PR, et al. [Predictors of poor mental and physical health status among patients with chronic hepatitis C infection: the chronic hepatitis cohort study \(CHeCS\)](#). *Hepatology*. 2015;61(3):802-811.
- Bourliere M, Gordon SC, Flamm SL, Cooper CL, Ramji A, Tong M, et al. [Sofosbuvir, velpatasvir, and voxilaprevir for previously treated HCV infection](#). *N Engl J Med*. 2017;376(22):2134-2146.
- Boursier J, de Ledinghen V, Zarski JP, Fouchard-Hubert I, Gallois Y, Oberti F, et al. [Comparison of eight diagnostic algorithms for liver fibrosis in hepatitis C: new algorithms are more precise and entirely noninvasive](#). *Hepatology*. 2012;55(1):58-67.
- Brau N, Salvatore M, Ríos-Bedoya CF, Fernandez-Carbia A, Paronetto F, Rodriguez-Orengo JF, et al. [Slower fibrosis progression in HIV/HCV-coinfected patients with successful HIV suppression using antiretroviral therapy](#). *J Hepatol*. 2006;44(1):47-55.
- Braun DL, Hampel B, Nguyen H. [A treatment as prevention trial to eliminate HCV in HIV+ MSM: the Swiss HCVfree trial \[abstract 81LB\]](#). 25th Conference on Retroviruses and Opportunistic Infections. Boston, MA; 2018.
- Bravo MJ, Vallejo F, Barrio G, Brugal MT, Molist G, Pulido J, et al. [HCV seroconversion among never-injecting heroin users at baseline: no predictors identified other than starting injection](#). *Int J Drug Policy*. 2012;23(5):415-419.
- Bruggmann P, Litwin AH. [Models of care for the management of hepatitis C virus among people who inject drugs: one size does not fit all](#). *Clin Infect Dis*. 2013;57(Suppl 2):S56-S61.
- Bruix J, Sherman M, AASLD. [Management of hepatocellular carcinoma: an update](#). *Hepatology*. 2011;53(3):1020-1022.
- Burton MJ, Voluse AC, Anthony V. [Integrating comprehensive hepatitis C virus care within a residential substance use disorder treatment program](#). *J Subst Abuse Treat*. 2019;(98):9-14.
- Butt AA, McGinnis K, Skanderson M, Justice AC. [A comparison of treatment eligibility for hepatitis C virus in HCV-monoinfected versus HCV/HIV-coinfected persons in electronically retrieved cohort of HCV-infected veterans](#). *AIDS Res Hum Retroviruses*. 2011;27(9):973-979.
- Canary LA, Klevens RM, Holmberg SD. [Limited access to new hepatitis C virus treatment under state medicaid programs](#). *Ann Intern Med*. 2015;163(3):226-228.
- Carrat F, Fontaine H, Dorival C. [Clinical outcomes in patients with chronic hepatitis C after direct-acting antiviral treatment: a prospective cohort study](#). *Lancet*. 2019;393(10179):1453-1464.
- Castera L, Vergniol J, Foucher J, Le BB, Chanteloup E, Haaser M, et al. [Prospective comparison of transient](#)

[elastography, Fibrotest, APRI, and liver biopsy for the assessment of fibrosis in chronic hepatitis C](#). Gastroenterology. 2005;128(2):343-350.

Castera L, Sebastiani G, Le BB, de Ledinghen V, Couzigou P, Alberti A. [Prospective comparison of two algorithms combining non-invasive methods for staging liver fibrosis in chronic hepatitis C](#). J Hepatol. 2010;52(2):191-198.

Castera L. [Noninvasive methods to assess liver disease in patients with hepatitis B or C](#). Gastroenterology. 2012;142(6):1293-1302.

[Centers for Disease Control and Prevention \(CDC\). Recommendations for prevention and control of hepatitis C virus \(HCV\) infection and HCV-related chronic disease](#). MMWR Recomm Rep. 1998;47(RR-19):1-39.

[Centers for Disease Control and Prevention \(CDC\). Recommendations for preventing transmission of infections among chronic hemodialysis patients](#). MMWR Recomm Rep. 2001;50(RR-5):1-43.

[Use of 13-valent pneumococcal conjugate vaccine and 23-valent pneumococcal polysaccharide vaccine for adults with immunocompromising conditions: recommendations of the Advisory Committee on Immunization Practices \(ACIP\)](#). MMWR Morb Mortal Wkly Rep. 2012;61(40):816-819.

[Centers for Disease Control and Prevention \(CDC\). Testing for HCV infection: an update of guidance for clinicians and laboratorians](#). MMWR Morb Mortal Wkly Rep. 2013;62(18):362-365.

Chahal HS, Marseille EA, Tice JA, Pearson SD, Ollendorf DA, Fox RK, et al. [Cost-effectiveness of early treatment of hepatitis C virus genotype 1 by stage of liver fibrosis in a US treatment-naive population](#). JAMA Intern Med. 2016;176(1):65-73.

Chaillon A, Rand EB, Reau N, Martin NK. [Cost-effectiveness of universal hepatitis C virus screening of pregnant women in the United States](#). Clin Infect Dis. 2019;69(11):1888-1895.

Chappell CA, Krans EE, Bunge K, et al. [A phase 1 study of ledipasvir/sofosbuvir in pregnant women with hepatitis C virus \[abstract 87\]](#). Presented at the Conference on Retroviruses and Opportunistic Infections; March 4-7. 2019.

Charlton MR, Seaberg E, Wiesner R, Everhart J, Zetterman R, Lake J, et al. [Predictors of patient and graft survival following liver transplantation for hepatitis C](#). Hepatology. 1998;28(3):823-830.

Charlton MR, Gane EJ, Manns MP, Brown, Jr. RS, Curry MP, Kwo PY, et al. [Sofosbuvir and ribavirin for treatment of compensated recurrent hepatitis C virus infection after liver transplantation](#). Gastroenterology. 2015;148(1):108-117.

Chen G, Wang C, Chen J. [Hepatitis B reactivation in hepatitis B and C coinfecting patients treated with antiviral agents: a systematic review and meta-analysis](#). Hepatology. 2017;66(1):13-26.

Chew KW, Allen SA, Taylor LE, Rich JD, Feller E. [Treatment outcomes with pegylated interferon and ribavirin for male prisoners with chronic hepatitis C](#). J Clin Gastroenterol. 2009;43(7):686-691.

Chhatwal J, Kanwal F, Roberts MS, Dunn MA. [Cost-effectiveness and budget impact of hepatitis C virus treatment with sofosbuvir and ledipasvir in the United States](#). Ann Intern Med. 2015;162(6):397-406.

- Chhatwal J, He T, Hur C, Lopez-Olivo MA. [Direct-acting antiviral agents for patients with hepatitis C virus genotype 1 infection are cost-saving](#). Clin Gastroenterol Hepatol. 2017;15(6):827-837.
- Chidi AP, Rogal S, Bryce CL, Bryce CL. [Cost-effectiveness of new antiviral regimens for treatment-naïve US veterans with hepatitis C](#). Hepatology. 2016;63(2):428-436.
- Chou R, Wasson N. [Blood tests to diagnose fibrosis or cirrhosis in patients with chronic hepatitis C virus infection: a systematic review](#). Ann Intern Med. 2013;158(11):807-820.
- Chou R, Dana T, Fu R, et al. [US Preventative Services Task Force. Screening for hepatitis C virus infection in adolescents and adults: updated evidence report and systematic review for the US Preventative Services Task Force](#). JAMA. 2020;.
- Chu CJ, Lee SD. [Hepatitis B virus/hepatitis C virus coinfection: epidemiology, clinical features, viral interactions and treatment](#). J Gastroenterol Hepatol. 2008;23(4):512-520.
- Clark BT, Garcia-Tsao G, Fraenkel L. [Patterns and predictors of treatment initiation and completion in patients with chronic hepatitis C virus infection](#). Patient Prefer Adherence. 2012;6:285-295.
- Comarmond C, Garrido M, Pol S. [Direct-acting antiviral therapy restores immune tolerance to patients with hepatitis C virus-induced cryoglobulinemia vasculitis](#). Gastroenterology. 2017;152(8):2052-2062.e2.
- Conjeevaram HS, Wahed AS, Afdhal NH, Howell CD, Everhart JE, Hoofnagle JH, et al. [Changes in insulin sensitivity and body weight during and after peginterferon and ribavirin therapy for hepatitis C](#). Gastroenterology. 2011;140(2):469-477.
- Cooper CL, Naggie S, Saag M, Yang JC, Stamm LM, Dvory-Sobol H, et al. [Successful re-treatment of hepatitis C virus in patients coinfecting with HIV who relapsed after 12 weeks of ledipasvir/sofosbuvir](#). Clin Infect Dis. 2016;63(4):528-531.
- Coppola N, De PS, Pisaturo M, Paradiso L, Macera M, Capoluongo N, et al. [Sustained virological response to antiviral treatment in chronic hepatitis C patients may be predictable by HCV-RNA clearance in peripheral blood mononuclear cells](#). J Clin Virol. 2013;58(4):748-750.
- Corrao G, Arico S. [Independent and combined action of hepatitis C virus infection and alcohol consumption on the risk of symptomatic liver cirrhosis](#). Hepatology. 1998;27(4):914-919.
- Coyle C, Viner K, Hughes E, Kwakwa H, Zibbell JE, Vellozzi C, et al. [Identification and linkage to care of HCV-infected persons in five health centers - Philadelphia, Pennsylvania, 2012-2014](#). MMWR Morb Mortal Wkly Rep. 2015;64(17):459-463.
- Curry MP, Forns X, Chung RT, Terrault NA, Brown, Jr. RS, Fenkel JM, et al. [Sofosbuvir and ribavirin prevent recurrence of HCV infection after liver transplantation: an open-label study](#). Gastroenterology. 2015;148(1):100-107.
- DeCarolis DD, Westanmo AD, Chen YC, Boese AL, Walquist MA, Rector TS. [Evaluation of a potential interaction between new regimens to treat hepatitis C and warfarin](#). Ann Pharmacother. 2016;(50):909-917.
- Denniston MM, Klevens RM, McQuillan GM, Jiles RB. [Awareness of infection, knowledge of hepatitis C, and medical](#)

[follow-up among individuals testing positive for hepatitis C: National Health and Nutrition Examination Survey 2001-2008.](#) Hepatology. 2012;55(6):1652-1661.

Di Bisceglie AM, Shiffman ML, Everson GT, Lindsay KL, Everhart JE, Wright EC, et al. [Prolonged therapy of advanced chronic hepatitis C with low-dose peginterferon.](#) N Engl J Med. 2008;359(23):2429-2441.

Dienstag JL, Ghany MG, Morgan TR, Di Bisceglie AM, Bonkovsky HL, Kim HY, et al. [A prospective study of the rate of progression in compensated, histologically advanced chronic hepatitis C.](#) Hepatology. 2011;54(2):396-405.

Dore GJ, Altice F, Litwin AH, Dalgard O, Gane EJ, Shibolet O, et al. [Elbasvir-grazoprevir to treat hepatitis C virus infection in persons receiving opioid agonist therapy: a randomized trial.](#) Ann Intern Med. 2016;165(9):625-634.

Drazilova S, Janicko M, Skladany L, et al. [Glucose metabolism changes in patients with chronic hepatitis C treated with direct acting antivirals.](#) Can J Gastroenterol Hepatol. 2018;2018:6095097.

Durier N, Nguyen C, White LJ. [Treatment of hepatitis C as prevention: a modeling case study in Vietnam.](#) PLoS One. 2012;7(4):e34548.

[European Association for the Study of the Liver; Asociacion Latinoamericana para el Estudio del Higado. EASL-ALEH clinical practice guidelines: non-invasive tests for evaluation of liver disease severity and prognosis.](#) J Hepatol. 2015;63(1):237-264.

Eckman MH, Ward JW, Sherman KE. [Cost effectiveness of universal screening for hepatitis C virus infection in the era of direct-acting, pangenotypic treatment regimens.](#) Clin Gastroenterol Hepatol. 2019;17(5):930-939.e9.

Edlin BR, Eckhardt BJ, Shu MA, Holmberg SD, Swan T. [Toward a more accurate estimate of the prevalence of hepatitis C in the United States.](#) Hepatology. 2015;62(5):1353-1363.

Emery JS, Kuczyński M, La D. [Efficacy and safety of direct acting antivirals for the treatment of mixed cryoglobulinemia.](#) Am J Gastroenterol. 2017;112(8):1298-1308.

Everhart JE, Lok AS, Kim HY, Morgan TR, Lindsay KL, Chung RT, et al. [Weight-related effects on disease progression in the hepatitis C antiviral long-term treatment against cirrhosis trial.](#) Gastroenterology. 2009;137(2):549-557.

Everhart JE, Wright EC, Goodman ZD, Dienstag JL, Hoefs JC, Kleiner DE, et al. [Prognostic value of Ishak fibrosis stage: findings from the hepatitis C antiviral long-term treatment against cirrhosis trial.](#) Hepatology. 2010;51(2):585-594.

Everson GT. [Treatment of patients with hepatitis C virus on the waiting list.](#) Liver Transpl. 2003;9(11):S90-S94.

Everson GT, Trotter J, Forman L, Kugelmas M, Halprin A, Fey B, et al. [Treatment of advanced hepatitis C with a low accelerating dosage regimen of antiviral therapy.](#) Hepatology. 2005;42(2):255-262.

Everson GT, Hoefs JC, Seeff LB, Bonkovsky HL, Naishadham D, Shiffman ML, et al. [Impact of disease severity on outcome of antiviral therapy for chronic hepatitis C: lessons from the HALT-C trial.](#) Hepatology. 2006;44(6):1675-1684.

Fabrizi F, Poordad F, Martin P. [Hepatitis C infection and the patient with end-stage renal disease.](#) Hepatology.

2002;36(1):3-10.

Fabrizi F, Takkouche B, Lunghi G, Dixit V, Messa P, Martin P. [The impact of hepatitis C virus infection on survival in dialysis patients: meta-analysis of observational studies](#). J Viral Hepat. 2007;14(10):697-703.

Fabrizi F, Messa P, Martin P. [Health-related quality of life in dialysis patients with HCV infection](#). Int J Artif Organs. 2009;32(8):473-481.

Fabrizi F, Dixit V, Messa P. [Antiviral therapy of symptomatic HCV-associated mixed cryoglobulinemia: meta-analysis of clinical studies](#). J Med Virol. 2013;85(6):1019-1027.

Fabrizi F, Martin P, Dixit V, Messa P. [Meta-analysis of observational studies: hepatitis C and survival after renal transplant](#). J Viral Hepat. 2014;21(5):314-324.

Feld JJ, Liang T.J. [Hepatitis C -- identifying patients with progressive liver injury](#). Hepatology. 2006;43(2 Suppl 1):S194-S206.

Fierer DS, Uriel AJ, Carriero DC, Klepper A, Dieterich D, Mullen MP, et al. [Liver fibrosis during an outbreak of acute hepatitis C virus infection in HIV-infected men: a prospective cohort study](#). J Infect Dis. 2008;198(5):683-686.

Finelli L, Miller JT, Tokars JI, Alter MJ, Arduino MJ. [National surveillance of dialysis-associated diseases in the United States, 2002](#). Semin Dial. 2005;18(1):52-61.

Fissell RB, Bragg-Gresham JL, Woods JD, Jadoul M, Gillespie B, Hedderwick SA, et al. [Patterns of hepatitis C prevalence and seroconversion in hemodialysis units from three continents: the DOPPS](#). Kidney Int. 2004;65(6):2335-2342.

Fontana RJ, Sanyal AJ, Ghany MG, Lee WM, Reid AE, Naishadham D, et al. [Factors that determine the development and progression of gastroesophageal varices in patients with chronic hepatitis C](#). Gastroenterology. 2010;138(7):2321-2331.

Ford MM, Jordan AE, Johnson N, Rude E, Laraque F, Varma JK, et al. [Check Hep C: a community-based approach to hepatitis C diagnosis and linkage to care in high-risk populations](#). J Public Health Manag Pract. 2018;24(1):41-48.

Forman LM, Lewis JD, Berlin JA, Feldman HI, Lucey MR. [The association between hepatitis C infection and survival after orthotopic liver transplantation](#). Gastroenterology. 2002;122(4):889-896.

Forns X, Navasa M, Rodes J. [Treatment of HCV infection in patients with advanced cirrhosis](#). Hepatology. 2004;40(2):498.

Forns X, Charlton MR, Denning J, McHutchison JG, Symonds WT, Brainard D, et al. [Sofosbuvir compassionate use program for patients with severe recurrent hepatitis C after liver transplantation](#). Hepatology. 2015;61(5):1485-1494.

Forns X, Lee SS, Valdes J, Lens S, Ghalib R, Aguilar H, et al. [Glecaprevir plus pibrentasvir for chronic hepatitis C virus genotype 1, 2, 4, 5, or 6 infection in adults with compensated cirrhosis \(EXPEDITION-1\): a single-arm, open-label, multicentre phase 3 trial](#). Lancet Infect Dis. 2017;17(10):1062-1068.

Foster GR, Goldin RD, Thomas HC. [Chronic hepatitis C virus infection causes a significant reduction in quality of life in the absence of cirrhosis](#). Hepatology. 1998;27(1):209-212.

Foster GR, Afdhal NH, Roberts SK. [Sofosbuvir and velpatasvir for HCV genotype 2 and 3 infection](#). N Engl J Med. 2015;373(27):2608-2617.

Foster GR, Gane E, Asatryan A, Asselah T, Ruane PJ, Pol S, et al. [ENDURANCE-3: safety and efficacy of glecaprevir/pibrentasvir compared to sofosbuvir plus daclatasvir in treatment-naive HCV genotype 3-infected patients without cirrhosis](#). J Hepatol [Internet]. 2017;66(1):S33. [http://dx.doi.org/10.1016/S0168-8278\(17\)30326-4](http://dx.doi.org/10.1016/S0168-8278(17)30326-4)

Gane EJ, Shiffman ML, Etzkorn K, Morelli G, Stedman C, Davis MN, et al. [Sofosbuvir/Velpatasvir in Combination With Ribavirin for 24 Weeks Is Effective Retreatment for Patients Who Failed Prior NS5A-Containing DAA Regimens: Results of the Retreatment Study](#). In 51st Annual Meeting of the European Association for the Study of the Liver (EASL). April 13-17. Barcelona, Spain; 2016.

Garcia-Bengoechea M, Basaras M, Barrio J, Arrese E, Montalvo II, Arenas JI, et al. [Late disappearance of hepatitis C virus RNA from peripheral blood mononuclear cells in patients with chronic hepatitis C in sustained response after alpha-interferon therapy](#). Am J Gastroenterol. 1999;94(7):1902-1905.

Garcia-Tsao G, Sanyal AJ, Grace ND, Carey W, Diseases practiceguidelines, Gastroenterology practiceparameters. [Prevention and management of gastroesophageal varices and variceal hemorrhage in cirrhosis](#). Hepatology. 2007;46(3):922-938.

Garcia-Tsao G, Abraldes JG, Berzigotti A, Bosch J. [Portal hypertensive bleeding in cirrhosis: Risk stratification, diagnosis, and management: 2016 practice guidance by the American Association for the Study of Liver Diseases](#). Hepatology. 2017;65(1):310-335.

George SL, Bacon BR, Brunt EM, Mihindukulasuriya KL, Hoffmann J, Di Bisceglie AM. [Clinical, virologic, histologic, and biochemical outcomes after successful HCV therapy: a 5-year follow-up of 150 patients](#). Hepatology. 2009;49(3):729-738.

Gerber L, Estep M, Stepanova M, Escheik C, Weinstein A, Younossi ZM. [Effects of viral eradication with ledipasvir and sofosbuvir, with or without ribavirin, on measures of fatigue in patients with chronic hepatitis C virus infection](#). Clin Gastroenterol Hepatol. 2016;14(1):156-164.e3.

Ghany MG, Kleiner DE, Alter H, Doo E, Khokar F, Promrat K, et al. [Progression of fibrosis in chronic hepatitis C](#). Gastroenterology. 2003;124(1):97-104.

Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R. [Prevalence of hepatitis C virus infection in porphyria cutanea tarda: systematic review and meta-analysis](#). J Hepatol. 2003;39(4):620-627.

Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R. [Systematic review: regression of lymphoproliferative disorders after treatment for hepatitis C infection](#). Aliment Pharmacol Ther. 2005;21(6):653-662.

Goel A, Chen Q, Chhatwal J, Aggarwal R. [Cost-effectiveness of generic pan-genotypic sofosbuvir/velpatasvir versus genotype-dependent direct-acting antivirals for hepatitis C treatment](#). J Gastroenterol Hepatol. 2018;33(12):2029-2036.

- Govindasamy D, Ford N, Kranzer K. [Risk factors, barriers and facilitators for linkage to antiretroviral therapy care: a systematic review](#). AIDS. 2012;26(16):2059-2067.
- Grady BP, Schinkel J, Thomas XV, Dalgard O. [Hepatitis C virus reinfection following treatment among people who use drugs](#). Clin Infect Dis. 2013;57(Suppl 2):S105-S110.
- Grebely J, Matthews GV, Hellard M, Shaw D, Shaw D, Petoumenos K, Yeung B, et al. [Adherence to treatment for recently acquired hepatitis C virus \(HCV\) infection among injecting drug users](#). J Hepatol. 2011;55(1):76-85.
- Gumber SC, Chopra S. [Hepatitis C: a multifaceted disease. Review of extrahepatic manifestations](#). Ann Intern Med. 1995;123(8):615-620.
- Hagström H. [Alcohol consumption in concomitant liver disease: how much is too much?](#). Curr Hepatol Rep. 2017;16(2):152-157.
- Harris DR, Gonin R, Alter HJ, Wright EC, Buskell ZJ, Hollinger FB, et al. [The relationship of acute transfusion-associated hepatitis to the development of cirrhosis in the presence of alcohol abuse](#). Ann Intern Med. 2001;134(2):120-124.
- Harris RJ, Martin NK, Rand E, Mandal S, Mutimer D, Vickerman P, et al. [New treatments for hepatitis C virus \(HCV\): scope for preventing liver disease and HCV transmission in England](#). J Viral Hepat. 2016;23(8):631-643.
- Harrison GI, Murray K, Gore R, et al. [The hepatitis C awareness through to treatment \(HepCATT\) study: improving the cascade of care for hepatitis C virus-infected people who inject drugs in England](#). Addiction. 2019;114(6):1113-1122.
- He T, Li K, Roberts MS, Spaulding AC, Ayer T, Grefenstette JJ, et al. [Prevention of hepatitis C by screening and treatment in US prisons](#). Ann Intern Med. 2016;164(2):84-92.
- He T, Lopez-Olivo MA, Hur C, Chhatwal J. [Systematic review: cost-effectiveness of direct-acting antivirals for treatment of hepatitis C genotypes 2-6](#). Aliment Pharmacol Ther. 2017;46(8):711-721.
- Hellard ME, Hocking JS, Crofts N. [The prevalence and the risk behaviours associated with the transmission of hepatitis C virus in Australian correctional facilities](#). Epidemiol Infect. 2004;132(3):409-15.
- Hellard ME, Jenkinson R, Higgs P, Stoope MA, Sacks-Davis R, Gold J, et al. [Modelling antiviral treatment to prevent hepatitis C infection among people who inject drugs in Victoria, Australia](#). Med J Aust. 2012;196(10):638-641.
- Hellard M, Doyle JS, Sacks-Davis R, Thompson AJ, McBryde E. [Eradication of hepatitis C infection: the importance of targeting people who inject drugs](#). Hepatology. 2014;59(2):366-369.
- Henderson DK, Dembry L, Fishman NO, Grady C, Lundstrom T, Palmore TN, et al. [SHEA guideline for management of healthcare workers who are infected with hepatitis B virus, hepatitis C virus, and/or human immunodeficiency virus](#). Infect Control Hosp Epidemiol. 2010;31(3):203-232.
- Hermine O, Lefrere F, Bronowicki JP, Mariette X, Jondeau K, Eclache-Saudreau V, et al. [Regression of splenic lymphoma with villous lymphocytes after treatment of hepatitis C virus infection](#). N Engl J Med. 2002;347(2):89-94.

- Hezode C, Reau N, Svarovskaia ES, Doehle BP, Shanmugam R, Dvory-Sobol H, et al. [Resistance analysis in patients with genotype 1-6 HCV infection treated with sofosbuvir/velpatasvir in the phase III studies](#). J Hepatol. 2018;68(5):895-903.
- Ho SB, Brau N, Cheung R, Liu L, Sanchez C, Sklar M, et al. [Integrated care increases treatment and improves outcomes of patients with chronic hepatitis C virus infection and psychiatric illness or substance abuse](#). Clin Gastroenterol Hepatol. 2015;13(11):2005-2014.e1-3.
- Hofmeister MG, Rosenthal EM, Barker LK, et al. [Estimating prevalence of hepatitis C virus infection in the United States, 2013-2016](#). Hepatology. 2019;69(3):1020-1031.
- Holmberg SD, Spradling PR, Moorman AC, Denniston MM. [Hepatitis C in the United States](#). N Engl J Med. 2013;368(20):1859-1861.
- Holmes HM, Hayley DC, Alexander GC, Sachs GA. [Reconsidering medication appropriateness for patients late in life](#). Arch Intern Med. 2006;166(6):605-609.
- Hoorneborg E, Coyer L, Boyd A, et al. [Amsterdam PrEP project team in the HIV transmission elimination Amsterdam \(H-TEAM\) initiative. High incidence of HCV in HIV-negative men who have sex with men using pre-exposure prophylaxis](#). J Hepatol. 2020;72(5):855-864.
- Hosein SR, Wilson DP. [HIV, HCV, and drug use in men who have sex with men](#). Lancet. 2013;382(9898):1095-1096.
- Hsieh YH, Rothman RE, Laeyendecker , et al. [Evaluation of the Centers for Disease Control and Prevention recommendations for hepatitis C virus testing in an urban emergency department](#). Clin Infect Dis. 2016;62(9):1059-1065.
- Hsu L, Bowlus CL, Stewart SL, Nguyen TT, Dang J, Chan B, et al. [Electronic messages increase hepatitis B screening in at-risk Asian American patients: a randomized, controlled trial](#). Dig Dis Sci. 2013;58(3):807-814.
- Hsu YC, Lin JT, Ho HJ, Kao YH, Huang YT, Hsiao NW, et al. [Antiviral treatment for hepatitis C virus infection is associated with improved renal and cardiovascular outcomes in diabetic patients](#). Hepatology. 2014;59(4):1293-1302.
- Hsu YC, Ho HJ, Huang YT, Wang HH, Wu MS, Lin JT, et al. [Association between antiviral treatment and extrahepatic outcomes in patients with hepatitis C virus infection](#). Gut. 2015;64(3):495-503.
- Hung CH, Wang JH, Hu TH, Chen CH, Chang KC, Yen YH, et al. [Insulin resistance is associated with hepatocellular carcinoma in chronic hepatitis C infection](#). World J Gastroenterol. 2010;16(18):2265-2271.
- Islam MM, Topp L, Conigrave KM, White A, Reid SE, Grummett S, et al. [Linkage into specialist hepatitis C treatment services of injecting drug users attending a needle syringe program-based primary healthcare centre](#). J Subst Abuse Treat. 2012;43(4):440-445.
- Jacobson IM, Asante-Appiah E, Wong P, Black T, Howe A, Wahl J, et al. [Prevalence and impact of baseline NSA resistance associated variants \(RAVs\) on the efficacy of elbasvir/grazoprevir \(EBR/GZR\) against GT1a infection \[abstract LB-22\]](#). The Liver Meeting. San Francisco, CA; 2015.

- Jadoul M, Cornu C, van Ypersele de Strihou C. [Universal precautions prevent hepatitis C virus transmission: a 54 month follow-up of the Belgian multicenter study. The Universitaires Cliniques St-Luc \(UCL\) collaborative group.](#) *Kidney Int.* 1998;53(4):1022-1025.
- Jezequel C, Bardou-Jacquet E, Desille Y, Renard I, Laine F, Lelan C, et al. [Survival of patients infected by chronic hepatitis C and F0F1 fibrosis at baseline after a 15 year follow-up \[P0709\].](#) *J Hepatol.* 2015;62(Suppl 2):S589.
- Johnson RJ, Gretch DR, Yamabe H, Hart J, Bacchi CE, Hartwell P, et al. [Membranoproliferative glomerulonephritis associated with hepatitis C virus infection.](#) *N Engl J Med.* 1993;328(7):465-470.
- Kanwal F, Kramer JR, Ilyas J, Duan Z, El-Serag HB. [HCV genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of US veterans with HCV.](#) *Hepatology.* 2014;60(1):98-105.
- Kattakuzhy S, Gross C, Emmanuel B. [Expansion of treatment for hepatitis C virus infection by task shifting to community-based nonspecialist providers: a nonrandomized clinical trial.](#) *Ann Intern Med.* 2017;167(5):311-318.
- [Kidney disease: improving global outcomes \(KDIGO\). KDIGO clinical practice guidelines for the prevention, diagnosis, evaluation, and treatment of hepatitis C in chronic kidney disease.](#) *Kidney Int Suppl.* 2008;(109):S1-S99.
- Khokhar OS, Lewis JH. [Reasons why patients infected with chronic hepatitis C virus choose to defer treatment: do they alter their decision with time?.](#) *Dig Dis Sci.* 2007;52(5):1168-1176.
- Konerman MA, Mehta SH, Sutcliffe CG, Vu T, Higgins Y, Torbenson MS, et al. [Fibrosis progression in human immunodeficiency virus/hepatitis C virus coinfecting adults: prospective analysis of 435 liver biopsy pairs.](#) *Hepatology.* 2014;59(3):767-775.
- Krishnan P, Pilot-Matias T, Schnell G, Tripathi R, Ng TI, Reisch T, et al. [Pooled resistance analysis in HCV genotype 1-6 infected patients treated with glecaprevir/pibrentasvir in phase 2 and 3 clinical trials.](#) *Antimicrob Agents Chemother.* 2018;62(10):pii: e01249-18.
- Kruse RL, Kramer JR, Tyson GL, Duan Z, Chen L, El-Serag HB, et al. [Clinical outcomes of hepatitis B virus coinfection in a United States cohort of hepatitis C virus-infected patients.](#) *Hepatology.* 2014;60(6):1871-1878.
- Kwo PY, Poordad F, Asatryan A, Wang S, Wyles DL, Hassanein T, et al. [Glecaprevir and pibrentasvir yield high response rates in patients with HCV genotype 1-6 without cirrhosis.](#) *J Hepatol.* 2017;67(2):263-271.
- Lai JC, Kahn JG, Tavakol M, Peters MG, Roberts JP. [Reducing infection transmission in solid organ transplantation through donor nucleic acid testing: a cost-effectiveness analysis.](#) *Am J Transplant.* 2013;13(10):2611-2618.
- Lambers FA, Prins M, Thomas X, et al, et al. [Alarming incidence of hepatitis C virus re-infection after treatment of sexually acquired acute hepatitis C virus infection in HIV-infected MSM.](#) *AIDS.* 2011;25(17):F21-F27.
- Landau DA, Scerra S, Sene D, Resche-Rigon M, Saadoun D, Cacoub P. [Causes and predictive factors of mortality in a cohort of patients with hepatitis C virus-related cryoglobulinemic vasculitis treated with antiviral therapy.](#) *J Rheumatol.* 2010;37(3):615-621.

- Lange B, Roberts T, Cohn J. [Diagnostic accuracy of detection and quantification of HBV-DNA and HCV-RNA using dried blood spot \(DBS\) samples - a systematic review and meta-analysis](#). BMC Infect Dis. 2017;17(Suppl 1):693.
- Larney S, Kopinski H, Beckwith CG, et al. [Incidence and prevalence of hepatitis C in prisons and other closed settings: results of a systematic review and meta-analysis](#). Hepatology. 2013;58(4):1215-1224.
- Lawitz EJ, Poordad F, Pang PS, Hyland RH, Ding X, Mo H, et al. [Sofosbuvir and ledipasvir fixed-dose combination with and without ribavirin in treatment-naive and previously treated patients with genotype 1 hepatitis C virus infection \(LONESTAR\): an open-label, randomised, phase 2 trial](#). Lancet. 2014;383(9916):515-523.
- Lee SR, Kardos KW, Schiff ER, Berne CA, Mounzer K, Banks AT, et al. [Evaluation of a new, rapid test for detecting HCV infection, suitable for use with blood or oral fluid](#). J Virol Methods. 2011;172(1-2):27-31.
- Lee SW, Lee TY, Yang SS, Peng YC, Yeh HZ, Chang CS. [Prevalence of hepatitis B reactivation among Chinese individuals with chronic hepatitis C treated with pan-oral direct-acting antivirals](#). Gastroenterology Res. 2018;11(2):124-129.
- Leroy V, Angus P, Bronowicki JP, et al. [Daclatasvir, sofosbuvir, and ribavirin for hepatitis C virus genotype 3 and advanced liver disease: A randomized phase III study \(ALLY-3+\)](#). Hepatology. 2016 Mar 4.th ed. 2016;63(5):1430-1441.
- Linac BP, Wong AY, Schackman BR, Kim AY, Freedberg KA. [Cost-effective screening for acute hepatitis C virus infection in HIV-infected men who have sex with men](#). Clin Infect Dis. 2012;55(2):279-290.
- Linac BP, Barter DM, Morgan JR, Pho MT, Leff JA, Schackman BR, et al. [The cost-effectiveness of sofosbuvir-based regimens for treatment of hepatitis C virus genotype 2 or 3 infection](#). Ann. Intern. Med. 2015;162(9):619-629.
- Litwin AH, Smith BD, Drainoni ML, McKee D, Gifford AL, Koppelman E, et al. [Primary care-based interventions are associated with increases in hepatitis C virus testing for patients at risk](#). Dig Liver Dis. 2012;44(6):497-503.
- Lo Re V, Kallan MJ, Tate JP, Localio AR, Lim JK, Goetz MB, et al. [Hepatic decompensation in antiretroviral-treated patients co-infected with HIV and hepatitis C virus compared with hepatitis C virus-monoinfected patients: a cohort study](#). Ann Intern Med. 2014;160(6):369-379.
- Lo Re V, Gowda C, Urlick PN, et al. [Disparities in absolute denial of modern hepatitis C therapy by type of insurance](#). Clin Gastroenterol Hepatol. 2016;14(7):1035-1043.
- Lok AS, McMahon BJ. [Chronic hepatitis B: update 2009](#). Hepatology. 2009;50(3):661-662.
- Louie KS, St Laurent S, Forssen UM, Mundy LM, Pimenta JM. [The high comorbidity burden of the hepatitis C virus infected population in the United States](#). BMC Infect Dis. 2012;12:86.
- Ly KN, Jiles RB, Teshale EH, Foster MA, Pesano RL, Holmberg SD. [Hepatitis C virus infection among reproductive-aged women and children in the United States, 2006 to 2014](#). Ann of Intern Med. 2017;166(11):775-782.
- Lyons MS, Kunnathur VA, Rouster SD, et al. [Prevalence of diagnosed and undiagnosed hepatitis C in a midwestern urban](#)

[emergency department](#). Clin Infect Dis. 2016;62(9):10-71.

Macias J, Berenguer J, Japon MA, et al. [Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus](#). Hepatology. 2009;50(4):1056-1063.

Maddison AR, Fisher J, Johnston G. [Preventive medication use among persons with limited life expectancy](#). Prog Palliat Care. 2011;19(1):15-21.

Manns MP, Pockros PJ, Norkrans G, Smith CI, Morgan TR, Haussinger D, et al. [Long-term clearance of hepatitis C virus following interferon alpha-2b or peginterferon alpha-2b, alone or in combination with ribavirin](#). J Viral Hepat. 2013;20(8):524-529.

Marcellin P, Boyer N, Gervais A, Martinot M, Pouteau M, Castelnau C, et al. [Long-term histologic improvement and loss of detectable intrahepatic HCV RNA in patients with chronic hepatitis C and sustained response to interferon-alpha therapy](#). Ann Intern Med. 1997;127(10):875-881.

Marrero JA, Kulik LM, Sirlin CB. [Diagnosis, staging, and management of hepatocellular carcinoma: 2018 practice guidance by the American Association for the Study of Liver Diseases](#). Hepatology. 2018;68(2):723-750.

Marrie TJ, Tyrrell GJ, Garg S, Vanderkooi O. [Factors predicting mortality in invasive pneumococcal disease in adults in Alberta](#). Medicine. 2011;90(3):171-179.

Martin NK, Hickman M, Hutchinson SJ, Goldberg DJ, Vickerman P. [Combination interventions to prevent HCV transmission among people who inject drugs: modeling the impact of antiviral treatment, needle and syringe programs, and opiate substitution therapy](#). Clin Infect Dis. 2013;57(Suppl 2):S39-S45.

Martin NK, Thornton A, Hickman M, et al. [Can hepatitis C virus \(HCV\) direct-acting antiviral treatment as prevention reverse the HCV epidemic among men who have sex with men in the United Kingdom? Epidemiological and modeling insights](#). Clin Infect Dis. 2016 Feb 16.th ed. 2016;62(9):1072-1080.

Martin NK, Vickerman P, Dore GJ, et al. [Prioritization of HCV treatment in the directacting antiviral era: an economic evaluation](#). J Hepatol. 2016;65(1):17-25.

Matheï C, Bourgeois S, Blach S, et al. [Mitigating the burden of hepatitis C virus among people who inject drugs in Belgium](#). Acta Gastroenterol Belg. 2016;79(2):227-232.

Mazzaro C, Little D, Pozzato G. [Regression of splenic lymphoma after treatment of hepatitis C virus infection](#). N Engl J Med. 2002;347(26):2168-2170.

McCombs JS, Tonnu-MiHara I, Matsuda T, McGinnis J, Fox S. [Can hepatitis C treatment be safely delayed? Evidence from the Veterans Administration healthcare system \[PIN104\]](#). Value Health. 2015;18(3):A245.

McGowan CE, Monis A, Bacon BR, Mallolas J, Goncales, Jr. FL, Goulis I, et al. [A global view of hepatitis C: physician knowledge, opinions, and perceived barriers to care](#). Hepatology. 2013;57(4):1325-1332.

Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL. [Prevalence of type 2 diabetes mellitus](#)

[among persons with hepatitis C virus infection in the United States](#). Ann Intern Med. 2000;133(8):592-599.

Mera J, Vellozzi C, Hariri S. [Identification and clinical management of persons with chronic hepatitis C virus infection - Cherokee Nation, 2012-2015](#). MMWR Morb Mortal Wkly Rep. 2016;65(18):461-466.

Merchante N, Giron-Gonzalez JA, Gonzalez-Serrano M, et al. [Survival and prognostic factors of HIV-infected patients with HCV-related end-stage liver disease](#). AIDS. 2006;20(1):49-57.

Midgard H, Bramness JG, Skurtveit S, Haukeland JW, Dalgard O. [Hepatitis C treatment uptake among patients who have received opioid substitution treatment: a population-based study](#). PloS One. 2016;11(11):e0166451.

Miller L, Fluker SA, Osborn M, Liu X, Strawder A. [Improving access to hepatitis C care for urban, underserved patients using a primary care-based hepatitis C clinic](#). J Natl Med Assoc. 2012;104(5-6):244-250.

Mira JA, Rivero-Juárez A, Lopez-Cortes LF, Giron-Gonzalez JA, Tellez F, Santos-Gil ID, et al. [Benefits from sustained virologic response to pegylated interferon plus ribavirin in HIV/hepatitis C virus-coinfected patients with compensated cirrhosis](#). Clin Infect Dis. 2013;56(11):1646-1653.

Morey S, Hamoodi A, Jones D, Young T. [Increased diagnosis and treatment of hepatitis C in prison by universal offer of testing and use of telemedicine](#). J Viral Hepat. 2019;26(1):101-108.

Morgan TR, Ghany MG, Kim HY, et al. [Outcome of sustained virological responders with histologically advanced chronic hepatitis C](#). Hepatology. 2010;52(3):833-844.

Morgan RL, Baack B, Smith BD, Yartel A, Pitasi M, Falck-Ytter Y. [Eradication of hepatitis C virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies](#). Ann Intern Med. 2013;158(5 Pt 1):329-337.

Morisco F, Granata R, Stroffolini T, Guarino M, Donnarumma L, Gaeta L, et al. [Sustained virological response: a milestone in the treatment of chronic hepatitis C](#). World J Gastroenterol. 2013;19(18):2793-2798.

Morrill JA, Shrestha M, Grant RW. [Barriers to the treatment of hepatitis C. Patient, provider, and system factors](#). J Gen Intern Med. 2005;20(8):754-758.

Moyer VA. [Screening for HIV: US Preventive Services Task Force recommendation statement](#). Ann Intern Med. 2013;159(1):51-60.

Mücke MM, Backus LI, Mücke VT. [Hepatitis B virus reactivation during direct-acting antiviral therapy for hepatitis C: a systematic review and meta-analysis](#). Lancet Gastroenterol Hepatol. 2018;3(3):172-180.

Murphy SM, Dweik D, McPherson S, Roll JM. [Association between hepatitis C virus and opioid use while in buprenorphine treatment: preliminary findings](#). Am J Drug Alcohol Abuse. 2015;41(1):88-92.

Najafzadeh M, Andersson K, Shrank WH, et al. [Cost-effectiveness of novel regimens for the treatment of hepatitis C virus](#). Ann Intern Med. 2015;162(6):407-419.

Neary MP, Cort S, Bayliss MS, Ware JE. [Sustained virologic response is associated with improved health-related quality of life in relapsed chronic hepatitis C patients](#). Semin Liver Dis. 1999;19(Suppl 1):77-85.

Nelson PK, Mathers BM, Cowie B, Hagan H, Des Jarlais D, Horyniak D, et al. [Global epidemiology of hepatitis B and hepatitis C in people who inject drugs: results of systematic reviews](#). Lancet. 2011;378(9791):571-583.

Nelson DR, Cooper JN, Lalezari JP, et al. [All-oral 12-week treatment with daclatasvir plus sofosbuvir in patients with hepatitis C virus genotype 3 infection: ALLY-3 phase III study](#). Lawitz EJ, Pockros PJ, Gitlin N, Freilich BF, Younes ZH, Harlan W, et al. Hepatology. 2015;61(4):1127-1135.

Neumann UP, Berg T, Bahra M, Seehofer D, Langrehr JM, Neuhaus R, et al. [Fibrosis progression after liver transplantation in patients with recurrent hepatitis C](#). J Hepatol. 2004;41(5):830-836.

Newsom AM, Stolte IG, van der Meer JTM. [Development and validation of the HCV-MOSAIC risk score to assist testing for acute hepatitis C virus \(HCV\) infection in HIV-infected men who have sex with men \(MSM\)](#). Euro Surveill. 2017;22(21):30540.

Ng T, Lu L, Reisch T, et al. [Resistance selection using glecaprevir and pibrentasvir in replicons of major hepatitis C virus genotypes](#). Krishnan P, Schnell G, Tripathi R, Beyer J, Dekhtyar T, Irvin M, et al. J Hepatol. 2017;66(1):S324.

Noda K, Yoshihara H, Suzuki K, Yamada Y, Kasahara A, Hayashi N, et al. [Progression of type C chronic hepatitis to liver cirrhosis and hepatocellular carcinoma--its relationship to alcohol drinking and the age of transfusion](#). Alcohol Clin Exp Res. 1996;20(1 Suppl):95A-100A.

Nutt AK, Hassan HA, Lindsey J, Lamps LW, Raufman JP. [Liver biopsy in the evaluation of patients with chronic hepatitis C who have repeatedly normal or near-normal serum alanine aminotransferase levels](#). Am J Med. 2000;109(1):62-64.

Owens DK, Davidson KW, Krist AH, et al. [US Preventive Services Task Force. Screening for hepatitis C virus infection in adolescents and adults: US Preventive Services Task Force recommendation statement](#). JAMA. 2020;.

Pakianathan M, Whittaker W, Lee M, et al. [Chemsex and new HIV diagnosis in gay, bisexual and other men who have sex with men attending sexual health clinics](#). HIV Med [Internet]. 2018;19(7):485-490. [10.1111/hiv.12629](#)

Pavone P, Tieghi T, d'Ettorre G, et al. [Rapid decline of fasting glucose in HCV diabetic patients treated with direct-acting antiviral agents](#). Clin Microbiol Infect. 2016;22(462):e461-463.

Pawlotsky JM. [Use and interpretation of virological tests for hepatitis C](#). Hepatology. 2002;36(5 Suppl 1):S65-S73.

Pawlotsky J-M. [Hepatitis C virus resistance to direct-acting antiviral drugs in interferon-free regimens](#). Gastroenterology. 2016;151(1):70-86.

Petta S, Camma C, di Marco V, et al. [Insulin resistance and diabetes increase fibrosis in the liver of patients with genotype 1 HCV infection](#). Alessi N, Cabibi D, Caldarella R, Licata A, Massenti F, Tarantino G, et al. Am J Gastroenterol. 2008;103(5):1136-1144.

Petta S, Maida M, Macaluso FS, et al. [Hepatitis C virus infection is associated with increased cardiovascular mortality: a](#)

[meta-analysis of observational studies](#). Barbara M, Licata A, Craxi A, Camma C. Gastroenterology. 2015 Sep 18th ed. 2016;150(1):145-155.e4.

Picciotto FP, Tritto G, Lanza AG, Addario L, De LM, Di Costanzo GG, et al. [Sustained virological response to antiviral therapy reduces mortality in HCV reinfection after liver transplantation](#). J Hepatol. 2007;46(3):459-465.

Pineda JA, Romero-Gomez M, Díaz-García F, et al. [HIV coinfection shortens the survival of patients with hepatitis C virus-related decompensated cirrhosis](#). Giron-Gonzalez JA, Montero JL, Torre-Cisneros J, Andrade RJ, Gonzalez-Serrano M, Aguilar J, et al. Hepatology. 2005;41(4):779-789.

Post JJ, Arain A, Lloyd AR. [Enhancing assessment and treatment of hepatitis C in the custodial setting](#). Clin Infect Dis. 2013;57(Suppl 2):S70-S74.

Poynard T, Bedossa P, Opolon P. [Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups](#). Lancet. 1997;349(9055):825-832.

Poynard T, Ratziu V, Charlotte F, Goodman Z, McHutchison JG, Albrecht J. [Rates and risk factors of liver fibrosis progression in patients with chronic hepatitis C](#). J Hepatol. 2001;34(5):730-739.

Poynard T, Cacoub P, Ratziu V, Myers RP, Dezailles MH, Mercadier A, et al. [Fatigue in patients with chronic hepatitis C](#). J Viral Hepat. 2002;9(4):295-303.

Poynard T, McHutchison JG, Manns M, Trepo C, Lindsay K, Goodman Z, et al. [Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C](#). Gastroenterology. 2002;122(5):1303-1313.

Pradat P, Alberti A, Poynard T, Esteban JI, Weiland O, Marcellin P, et al. [Predictive value of ALT levels for histologic findings in chronic hepatitis C: a European collaborative study](#). Hepatology. 2002;36(4 Pt 1):973-977.

Puoti M, Lorenzini P, Cozzi-Lepri A, et al. [Incidence and progression to cirrhosis of new hepatitis C virus infections in persons living with human immunodeficiency virus](#). Gori A, Mastroianni C, Rizzardini G, Mazzarello G, Antinori A, Monforte Ad'Arminio, et al. Clin Microbiol Infect. 2017;23(4):267.e1-267.e4.

Reddy KR, Pol S, Thuluvath PJ. [Long-term follow-up of clinical trial patients treated for chronic HCV infection with daclatasvir-based regimens](#). Liver Int. 2018;38(5):821-833.

Reig M, Mariño Z, Perelló C, et al. [Unexpected high rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free therapy](#). Iñarrairaegui M, Ribeiro A, Lens S, Díaz A, Vilana R, Darnell A, et al. J Hepatol. 2016;65(4):719-726.

Reilley B, Leston J, Redd JT, Geiger R. [Lack of access to treatment as a barrier to HCV screening: a facility-based assessment in the Indian Health Service](#). J Public Health Manag Pract. 2014;20(4):420-423.

Rein DB, Wittenborn JS, Smith BD, et al. [The cost-effectiveness, health benefits, and financial costs of new antiviral treatments for hepatitis C virus](#). Clin Infect Dis. 2015;61(2):157-168.

Rindone JP, Mellen CK. [Reduction in warfarin effect associated with sofosbuvir-velpatasvir](#). Am J Health Syst Pharm.

2017;(74):1308-1311.

Rockey DC, Bissell DM. [Noninvasive measures of liver fibrosis](#). Hepatology. 2006;43(2 Suppl 1):S113-S120.

Rossaro L, Torruellas C, Dhaliwal S, et al. [Clinical outcomes of hepatitis C treated with pegylated interferon and ribavirin via telemedicine consultation in Northern California](#). Botros J, Clark G, Li CS, Minoletti MM. Dig Dis Sci. 2013;58(12):3620-3625.

Saag MS. [Editorial commentary: getting smart in how we pay for HCV drugs: KAOS vs CONTROL](#). Clin Infect Dis. 2015;61(2):169-170.

Safdar K, Schiff ER. [Alcohol and hepatitis C](#). Semin Liver Dis. 2004;24(3):305-315.

Samandari T, Tedaldi E, Armon C, et al. [Incidence of hepatitis C virus infection in the human immunodeficiency virus outpatient study cohort, 2000-2013](#). Hart R, Chmiel JS, Brooks JT, Buchacz K, Investigators and the HIVOU. Open Forum Infect Dis. 2017;4(2):ofx076.

Sangiovanni A, Prati GM, Fasani P, Ronchi G, Romeo R, Manini M, et al. [The natural history of compensated cirrhosis due to hepatitis C virus: a 17-year cohort study of 214 patients](#). Hepatology. 2006;43(6):1303-1310.

Sarkar S, Jiang Z, Evon DM, Wahed AS, Hoofnagle JH. [Fatigue before, during and after antiviral therapy of chronic hepatitis C: results from the Virahep-C study](#). J Hepatol. 2012;57(5):946-952.

Sarrazin C, Dvory-Sobol H, Svarovskaia ES, et al. [Prevalence of resistance-associated substitutions in HCV NS5A, NS5B, or NS3 and outcomes of treatment with ledipasvir and sofosbuvir](#). Doehle BP, Pang PS, Chuang SM, Ma J, Ding X, Afdhal NH, et al. Gastroenterology. 2016 June 11th ed. 2016;151(3):501-512.E1.

Sarrazin C, Isakov V, Svarovskaia ES. [Late relapse versus hepatitis C virus reinfection in patients with sustained virologic response after sofosbuvir-based therapies](#). Clin Infect Dis. 2017;64(1):44-52.

Sayiner M, Golabi P, Farhat F, Younossi ZM. [Dermatologic manifestations of chronic hepatitis C infection](#). Clin Liver Dis. 2017;21(3):555-564.

Schackman BR, Leff JA, Barter DM, et al. [Cost-effectiveness of rapid hepatitis C virus \(HCV\) testing and simultaneous rapid HCV and HIV testing in substance abuse treatment programs](#). Addiction. 2015;110(1):129-143.

Schackman BR, Gutkind S, Morgan JR, et al. [Cost-effectiveness of hepatitis C screening and treatment linkage intervention in US methadone maintenance treatment programs](#). Drug Alcohol Depend. 2018;(185):411-420.

Schechter-Perkins EM, Miller NS, Hall J, et al. [Implementation and preliminary results of an emergency department nontargeted, opt-out hepatitis C virus screening program](#). Acad Emerg Med. 2018;25(11):1226.

Schillie S, Wester C, Osborne M, Wesolowski L, Ryerson AB. [CDC recommendations for hepatitis C screening among adults - United States, 2020](#). MMWR Recomm Rep. 2020;69(2):1-17.

Schneider MD, Sarrazin C. [Antiviral therapy of hepatitis C in 2014: do we need resistance testing?](#) Antiviral Res.

2014;105:64-71.

Schulkind J, Stephens B, Ahmad F, et al. [High response and re-infection rates among people who inject drugs treated for hepatitis C in a community needle and syringe programme](#). J Viral Hepat. 2019;26(5):519-528.

Sebastiani G, Halfon P, Castera L, Pol S, Thomas DL, Mangia A, et al. [SAFE biopsy: a validated method for large-scale staging of liver fibrosis in chronic hepatitis C](#). Hepatology. 2009;49(6):1821-1827.

Seem DL, Lee I, Umscheid CA, Kuehnert MJ. [Excerpt from PHS guideline for reducing HIV, HBV and HCV transmission through organ transplantation](#). Am J Transplant. 2013;13(8):1953-1962.

Selph S, Chou R. [Impact of contacting study authors on systematic review conclusions: diagnostic tests for hepatic fibrosis](#). 2014.

Simmons B, Saleem J, Hill A, Riley RD, Cooke GS. [Risk of late relapse or reinfection with hepatitis C virus after achieving a sustained virological response: a systematic review and meta-analysis](#). Clin Infect Dis. 2016;62(6):683-694.

Singal AG, Volk ML, Jensen D, Di Bisceglie AM, Schoenfeld PS. [A sustained viral response is associated with reduced liver-related morbidity and mortality in patients with hepatitis C virus](#). Clin Gastroenterol Hepatol. 2010;8(3):280-288, 288e1.

Sise ME, Bloom AK, Wisocky J, et al. [Treatment of hepatitis C virus-associated mixed cryoglobulinemia with direct-acting antiviral agents](#). Lin MV, Gustafson JL, Lundquist AL, Steele D, Thiim M, Williams WW, et al. Hepatology. 2016;63(2):408-417.

Smith BD, Morgan RL, Beckett GA, et al, et al. [Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945-1965](#). Falck-Ytter Y, Holtzman D, Teo CG, Jewett A, Baack B, Rein DB, et al. MMWR Recomm Rep. 2012;61(RR-4):1-32.

Soriano V, Barreiro P, deMendoza C. [Hypoglycemia in a diabetic patient during hepatitis C therapy](#). Hepatology. 2016;(63):2065-2066.

Spaulding AC, Weinbaum CM, Lau DT, et al. [A framework for management of hepatitis C in prisons](#). Sterling RK, Seeff LB, Margolis HS, Hoofnagle JH. Ann Intern Med. 2006;144(10):762-769.

Stein MR, Soloway IJ, Jefferson KS, Roose RJ, Arnsten JH, Litwin AH. [Concurrent group treatment for hepatitis C: implementation and outcomes in a methadone maintenance treatment program](#). J Subst Abuse Treat. 2012;43(4):424-432.

Suryaprasad AG, White JZ, Xu F. [Emerging epidemic of hepatitis C virus infections among young nonurban persons who inject drugs in the United States, 2006-2012](#). Clin Infect Dis. 2014;(59):1411-1419.

Svarovskaia ES, Dvory-Sobol H, Parkin N, et al. [Infrequent development of resistance in genotype 1-6 hepatitis C virus-infected subjects treated with sofosbuvir in phase 2 and 3 clinical trials](#). Hebner C, Gontcharova V, Martin R, Ouyang W, Han B, Xu S, et al. Clin Infect Dis. 2014;59(12):1666-1674.

- Svoboda J, Andreadis C, Downs LH, Miller WT, Tsai DE, Schuster SJ. [Regression of advanced non-splenic marginal zone lymphoma after treatment of hepatitis C virus infection](#). *Leuk Lymphoma*. 2005;46(9):1365-1368.
- Swain MG, Lai MY, Shiffman ML, Cooksley WG, Zeuzem S, Dieterich D, et al. [A sustained virologic response is durable in patients with chronic hepatitis C treated with peginterferon alfa-2a and ribavirin](#). *Gastroenterology*. 2010;139(5):1593-1601.
- Takahashi K, Nishida N, Kawabata H, Haga H, Chiba T. [Regression of Hodgkin lymphoma in response to antiviral therapy for hepatitis C virus infection](#). *Intern Med*. 2012;51(19):2745-2747.
- Takikawa H, Yamazaki R, Shoji S, Miyake K, Yamanaka M. [Normalization of urinary porphyrin level and disappearance of skin lesions after successful interferon therapy in a case of chronic hepatitis C complicated with porphyria cutanea tarda](#). *J Hepatol*. 1995;22(2):249-250.
- Talal AH, Andrews P, Mcleod A, et al. [Integrated, co-located, telemedicine-based treatment approaches for hepatitis C virus \(HCV\) management in opioid use disorder patients on methadone](#). *Clin Infect Dis*. 2019;69(2):323-331.
- Tasillo A, Eftekhari-Yazdi G, Nolen S, et al. [Short-term effects and long-term cost-effectiveness of universal hepatitis C testing in prenatal care](#). *Obstet Gynecol*. 2019;133(2):289-300.
- Terrault NA, Roland ME, Schiano T, Dove L, Wong MT, Poordad F, et al. [Outcomes of liver transplant recipients with hepatitis C and human immunodeficiency virus coinfection](#). *Liver Transpl*. 2012;18(6):716-726.
- Terrault NA, Lok ASF, McMahon BJ, et al. [Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance](#). *Hepatology*. 2018;67(4):1560-1599.
- Thein HH, Yi Q, Dore GJ, Krahn MD. [Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis](#). *AIDS*. 2008;22(15):1979-1991.
- Thomas DL, Villano SA, Riester KA, Hershow R, Mofenson LM, Landesman SH, et al. [Perinatal transmission of hepatitis C virus from human immunodeficiency virus type 1-infected mothers. Women and infants transmission study](#). *J Infect Dis*. 1998;177(6):1480-1488.
- Tice JA, Chahal HS, Ollendorf DA. [Comparative clinical effectiveness and value of novel interferon-free combination therapy for hepatitis C genotype 1 summary of California technology assessment forum report](#). *JAMA Intern Med*. 2015;175(9):1559-1560.
- Torres HA, Mahale P. [Most patients with HCV-associated lymphoma present with mild liver disease: a call to revise antiviral treatment prioritization](#). *Liver Int*. 2015;35(6):1661-1664.
- Torres HA, Hosry J, Mahale P, Economides MP, Jiang Y, Lok AS. [Hepatitis C virus reactivation in patients receiving cancer treatment: A prospective observational study](#). *Hepatology*. 2018;67(1):36-47.
- Trooskin SB, Poceta J, Towey CM, Yolken A, Rose JS, Luqman NL, et al. [Results from a geographically focused, community-based HCV screening, linkage-to-care and patient navigation program](#). *J Gen Intern Med*. 2015;30(7):950-957.

Tyson GL, Kramer JR, Duan Z, Davila JA, Richardson PA, El-Serag HB. [Prevalence and predictors of hepatitis B virus coinfection in a United States cohort of hepatitis C virus-infected patients](#). Hepatology. 2013;58(2):538-545.

Urbanus AT, van de Laar TJ, Stolte IG, Schinkel J, Heijman T, Coutinho RA, et al. [Hepatitis C virus infections among HIV-infected men who have sex with men: an expanding epidemic](#). AIDS. 2009;23(12):F1-F7.

[Prescription drug pricing in the private sector](#). 2015.

US GAO. [An overview of approaches to negotiate drug prices used by other countries and US private payers and federal programs](#) [Internet]. 2015. <http://www.gao.gov/products/GAO-07-358T>

US Senate Committee on Finance. [The price of Sovaldi and its impact on the US health care system](#). 2016.

van de Laar T, Pybus O, Bruisten S, Brown D, Nelson M, Bhagani S, et al. [Evidence of a large, international network of HCV transmission in HIV-positive men who have sex with men](#). Gastroenterology. 2009;136(5):1609-1617.

van de Laar TJ, Matthews GV, Prins M, Danta M. [Acute hepatitis C in HIV-infected men who have sex with men: an emerging sexually transmitted infection](#). AIDS. 2010;24(12):1799-1812.

van der Meer AJ, Veldt BJ, Feld JJ, Wedemeyer H, Dufour JF, Lammert F, et al. [Association between sustained virological response and all-cause mortality among patients with chronic hepatitis C and advanced hepatic fibrosis](#). JAMA. 2012;308(24):2584-2593.

vanTilborg M, Al-Marzooqi SH, Wong WWL. [HCV core antigen as an alternative to HCV RNA testing in the era of direct-acting antivirals: retrospective screening and diagnostic cohort studies](#). Lancet Gastroenterol Hepatol. 2018;3(12):856-864.

Veldt BJ, Heathcote EJ, Wedemeyer H, Reichen J, Hofmann WP, Zeuzem S, et al. [Sustained virologic response and clinical outcomes in patients with chronic hepatitis C and advanced fibrosis](#). Ann Intern Med. 2007;147(10):677-684.

Vermeersch P, Van RM, Lagrou K. [Validation of a strategy for HCV antibody testing with two enzyme immunoassays in a routine clinical laboratory](#). J Clin Virol. 2008;42(4):394-398.

Volk JE, Marcus JL, Phengrasamy T, et al. [No new HIV Infections with increasing use of HIV preexposure prophylaxis in a clinical practice setting](#). Clin Infect Dis. 2015;61(10):1601-1603.

Wai CT, Greenson JK, Fontana RJ, Kalbfleisch JD, Marrero JA, Conjeevaram HS, et al. [A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C](#). Hepatology. 2003;38(2):518-526.

Wandeler G, Gsponer T, Bregenzer A, Gunthard HF, Clerc O, Calmy A, et al. [Hepatitis C virus infections in the Swiss HIV cohort study: a rapidly evolving epidemic](#). Clin Infect Dis. 2012;55(10):1408-1416.

Ward KM, Falade-Nwulia O, Moon J. [Randomized controlled trial of cash incentives or peer mentors to improve HCV linkage and treatment among HIV/HCV coinfecting persons who inject drugs: the CHAMPS Study](#). Open Forum Infect Dis. 2019;6(4):ofz166.

- Waziry R, Hajarizadeh B, Grebely J, et al. [Hepatocellular carcinoma risk following direct-acting antiviral HCV therapy: a systematic review, meta-analyses, and meta-regression](#). J Hepatol. 2017;67(6):1204-1212.
- Wedemeyer H, Duberg AS, Buti M, Rosenberg WM, Frankova S, Esmat GE, et al. [Strategies to manage hepatitis C virus \(HCV\) disease burden](#). J Viral Hepat. 2014;21(Suppl 1):60-89.
- Weinbaum C, Lyster R, Margolis HS, Margolis HS. [Prevention and control of infections with hepatitis viruses in correctional settings](#). MMWR Recomm Rep. 2003;52(RR-1):1-36.
- Weinbaum CM, Williams IT, Mast EE, Mast EE. [Centers for Disease Control and Prevention \(CDC\). Recommendations for identification and public health management of persons with chronic hepatitis B virus infection](#). MMWR Recomm Rep. 2008;57(RR-8):1-20.
- Westin J, Lagging LM, Spak F, Aires N, Svensson E, Lindh M, et al. [Moderate alcohol intake increases fibrosis progression in untreated patients with hepatitis C virus infection](#). J Viral Hepat. 2002;9(3):235-241.
- White DL, Ratziu V, El-Serag HB. [Hepatitis C infection and risk of diabetes: a systematic review and meta-analysis](#). J Hepatol. 2008;49(5):831-844.
- Wiley TE, McCarthy M, Breidi L, McCarthy M, Layden TJ. [Impact of alcohol on the histological and clinical progression of hepatitis C infection](#). Hepatology. 1998;28(3):805-809.
- Williams IT, Bell BP, Kuhnert W, Alter MJ. [Incidence and transmission patterns of acute hepatitis C in the United States, 1982-2006](#). Arch Intern Med. 2011;171(3):242-248.
- Witt MD, Seaberg EC, Darilay A, Young S, Badri S, Rinaldo CR, et al. [Incident hepatitis C virus infection in men who have sex with men: a prospective cohort analysis, 1984-2011](#). Clin Infect Dis. 2013;57(1):77-84.
- Wohl DA, Allmon AG, Evon D. [Financial incentives for adherence to hepatitis C virus clinical care and treatment: a randomized trial of two strategies](#). Open Forum Infect Dis. 2017;4(2):ofx095.
- Wong JB, Cohen JT. [Cost-effective but bad for health? Hepatitis C treatment, moral hazard, and opportunity cost](#). Clin Gastroenterol Hepatol. 2017;15(6):838-840.
- Wyles D, Mangia A, Cheng W, Shafran S, Schwabe C, Ouyang W, et al. [Long-term persistence of HCV NS5A resistance associated substitutions after treatment with the HCV NS5A inhibitor, ledipasvir, without sofosbuvir \[Epub ahead of print\]](#). Antiviral Therapy. 2017;.
- Yoneda M, Saito S, Ikeda T, Fujita K, Mawatari H, Kirikoshi H, et al. [Hepatitis C virus directly associates with insulin resistance independent of the visceral fat area in nonobese and nondiabetic patients](#). J Viral Hepat. 2007;14(9):600-607.
- Younossi ZM, Zheng L, Stepanova M, Venkatesan C, Mir HM. [Moderate, excessive or heavy alcohol consumption: each is significantly associated with increased mortality in patients with chronic hepatitis C](#). Aliment Pharmacol Ther. 2013;37(7):703-709.

- Younossi ZM, Stepanova M, Nader F, Jacobson IM, Gane EJ, Nelson DR, et al. [Patient-reported outcomes in chronic hepatitis C patients with cirrhosis treated with sofosbuvir-containing regimens](#). Hepatology. 2014;59(6):2161-2169.
- Younossi ZM, Stepanova M, Henry L, Gane EJ, Jacobson IM, Lawitz EJ, et al. [Effects of sofosbuvir-based treatment, with and without interferon, on outcome and productivity of patients with chronic hepatitis C](#). Clin Gastroenterol Hepatol. 2014;12(8):1349-1359.
- Younossi ZM, Park H, Saab S, Ahmed A, Dieterich D, Gordon SC. [Cost-effectiveness of all-oral ledipasvir/sofosbuvir regimens in patients with chronic hepatitis C virus genotype 1 infection](#). Aliment. Pharmacol. Ther. 2015;41(6):544-563.
- Younossi ZM, Jiang Y, Smith NJ, Stepanova M, Beckerman R. [Ledipasvir/sofosbuvir regimens for chronic hepatitis C infection: Insights from a work productivity economic model from the United States](#). Hepatology. 2015;61(5):1471-1478.
- Younossi ZM, Stepanova M, Afdhal NH, Kowdley KV, Zeuzem S, Henry L, et al. [Improvement of health-related quality of life and work productivity in chronic hepatitis C patients with early and advanced fibrosis treated with ledipasvir and sofosbuvir](#). J Hepatol. 2015;63(2):337-345.
- Younossi ZM, Stepanova M, Sulkowski MS, Naggie S, Puoti M, Orkin C, et al. [Sofosbuvir and ribavirin for treatment of chronic hepatitis C in patients coinfecting with hepatitis C virus and HIV: the impact on patient-reported outcomes](#). J Infect Dis. 2015;212(3):367-377.
- Younossi ZM, Stepanova M, Feld J, Zeuzem S, Jacobson I, Agarwal K, et al. [Sofosbuvir/velpatasvir improves patient-reported outcomes in HCV patients: results from ASTRAL-1 placebo-controlled trial](#). J Hepatol. 2016;65(1):33-39.
- Younossi ZM, Stepanova M, Henry L, Nader F, Hunt S. [An in-depth analysis of patient-reported outcomes in patients with chronic hepatitis C treated with different anti-viral regimens](#). Am J Gastroenterol. 2016;111(6):808-816.
- Younossi ZM, Stepanova M, Jacobson I. [Not achieving sustained viral eradication of hepatitis C virus after treatment leads to worsening of patient-reported outcomes](#). Clin Infect Dis. 2019;pii: ciz243.
- Zahnd C, Salazar-Vizcaya LP, Dufour JF, Mullhaupt B, Wandeler G, Kouyos R, et al. [Impact of deferring HCV treatment on liver-related events in HIV+ patients](#). In Conference on Retroviruses and Opportunistic Infections (CROI) February 23-26. Seattle, WA; 2015.
- Zarski JP, Bohn B, Bastie A, Pawlotsky JM, Baud M, Bost-Bezeaux F, et al. [Characteristics of patients with dual infection by hepatitis B and C viruses](#). J Hepatol. 1998;28(1):27-33.
- [Merck & Co. Inc. Zepatier Package insert](#). Merck Sharp & Dohme Corp; 2019.
- Zeuzem S, Ghalib R, K. Reddy R, Pockros PJ, Ben Ari, iv Z, Zhao Y, et al. [Grazoprevir-elbasvir combination therapy for treatment-naive cirrhotic and noncirrhotic patients with chronic hepatitis C virus genotype 1, 4, or 6 infection: a randomized trial](#). Ann Intern Med. 2015;163(1):1-13.
- Zeuzem S, Feld J, Wang S. [ENDURANCE-1: efficacy and safety of 8- versus 12-week treatment with ABT-493/ABT-530 in patients with chronic HCV genotype 1 infection \[abstract 253\]](#). In The Liver Meeting 2016. Boston, MA; 2016.

Zeuzem S, Mizokami M, Pianko S, Mangia A, Han KH, Martin R, et al. [NS5A resistance-associated substitutions in patients with genotype 1 hepatitis C virus: prevalence and effect on treatment outcome](#). J Hepatol. 2017;66(5):910-918.

Zibbell JE, Asher AK, Patel RC. [Increases in acute hepatitis C virus infection related to a growing opioid epidemic and associated injection drug use, United States, 2004 to 2014](#). Am J Public Health. 2018;108(2):175-181.

Ziol M, Handra-Luca A, Kettaneh A, Christidis C, Mal F, Kazemi F, et al. [Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with chronic hepatitis C](#). Hepatology. 2005;41(1):48-54.

Øvrehus ALH, Blach S, Christensen PB, Gerstoft J, Weis N, Krarup H, et al.. [Impact of prioritizing treatment in a high resource setting - minimizing the burden of HCV related disease in 15 years \[P074\]](#). J Hepatol. 2015;62(Suppl 2):S591-S592.