




## Treatment-Naive Genotype 1a Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for: <b>Treatment-Naive Genotype 1a Patients Without Cirrhosis</b>		
RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>a</sup>	8 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks	I, B
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
ALTERNATIVE	DURATION	RATING
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
<sup>a</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.		


**Last update:** October 24, 2022

## Treatment-Naive Genotype 1a With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for: <b>Treatment-Naive Genotype 1a Patients With Compensated Cirrhosis<sup>a</sup></b> 		
RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks	I, B

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Genotype 1a Patients With Compensated Cirrhosis<sup>a</sup>


Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
<sup>a</sup> For <a href="#">decompensated cirrhosis</a> , please refer to the appropriate section. <sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.		

**Last update:** October 24, 2022

## Treatment-Naive Genotype 1b Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Patients Genotype 1b Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>a</sup>	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks <sup>b</sup>	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for patients who are HIV-uninfected and whose HCV RNA level is <6 million IU/mL	8 weeks <sup>c</sup>	I, B
<sup>a</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information. <sup>b</sup> For HIV/HCV coinfecting patients, a treatment duration of 12 weeks is recommended. <sup>c</sup> An 8-week regimen can be considered in those with genotype 1b infection and mild fibrosis (see text for details).		

**Last update:** October 24, 2022

## Treatment-Naive Genotype 1b With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Genotype 1b Patients With Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks	I, B
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	I, A

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.

<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

**Last update:** October 24, 2022

## Treatment-Naive Genotype 2 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Genotype 2 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>a</sup>	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A


<sup>a</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

**Last update:** October 24, 2022

## Treatment-Naive Genotype 2 With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Genotype 2 Patients With Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks	I, B

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.


<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

**Last update:** October 24, 2022

## Treatment-Naive Genotype 3 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:




### Treatment-Naive Genotype 3 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>a</sup>	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A

<sup>a</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

**Last update:** October 24, 2022

## Treatment-Naive Genotype 3 With Compensated Cirrhosis


Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for: <b>Treatment-Naive Genotype 3 Patients With Compensated Cirrhosis<sup>a</sup></b> 		
RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) for patients without baseline NS5A RAS Y93H	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks	I, B
ALTERNATIVE	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg) with weight-based ribavirin for patients with baseline NS5A RAS Y93H	12 weeks	IIa, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)/voxilaprevir (100 mg) for patients with baseline NS5A RAS Y93H	12 weeks	IIa, B
<sup>a</sup> For <a href="#">decompensated cirrhosis</a> , please refer to the appropriate section. <sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.		

**Last update:** October 24, 2022

## Treatment-Naive Genotype 4 Without Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

## Treatment-Naive Genotype 4 Patients Without Cirrhosis

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>a</sup>	8 weeks	I, A
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) <sup>b</sup>	12 weeks	I, A

<sup>a</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.


<sup>b</sup> An 8-week regimen can be considered in patients with favorable baseline characteristics (ie, no cirrhosis, HCV RNA <6 million IU/mL, and absence of genotype 4r).

**Last update:** October 24, 2022

## Treatment-Naive Genotype 4 With Compensated Cirrhosis

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

## Treatment-Naive Genotype 4 Patients With Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, A
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks <sup>c</sup>	I, B
Daily fixed-dose combination of elbasvir (50 mg)/grazoprevir (100 mg)	12 weeks	IIa, B
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg)	12 weeks	IIa, B

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.

<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.


<sup>c</sup> For HIV/HCV-coinfected patients, a treatment duration of 12 weeks is recommended.

**Last update:** October 24, 2022

## Treatment-Naive Genotype 5 or 6

Recommended and alternative regimens listed by pangenotypic, evidence level and alphabetically for:

### Treatment-Naive Genotype 5 or 6 Patients With and Without Compensated Cirrhosis<sup>a</sup>

RECOMMENDED	DURATION	RATING 
Daily fixed-dose combination of glecaprevir (300 mg)/pibrentasvir (120 mg) <sup>b</sup>	8 weeks	I, A <sup>c</sup>
Daily fixed-dose combination of sofosbuvir (400 mg)/velpatasvir (100 mg)	12 weeks	I, B
Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) <sup>d</sup>	12 weeks	IIa, B

<sup>a</sup> For [decompensated cirrhosis](#), please refer to the appropriate section.

<sup>b</sup> Dosing is 3 coformulated tablets (glecaprevir [100 mg]/pibrentasvir [40 mg]) taken once daily. Please refer to the prescribing information.

<sup>c</sup> For compensated cirrhosis, rating is I, B.

<sup>d</sup> Not recommended for genotype 6e if subtype is known.

**Last update:** October 24, 2022